

Three Dollars per Annum, in Advance. Single Copies, Twenty-Five Cents.

VOL. II.

No. IV.

THE

# MATHEMATICAL MONTHLY.

JANUARY, 1860.

EDITED BY

J. D. RUNKLE, A.M., A.A.S.

NEW YORK:

PUBLISHED BY IVISON, PHINNEY, & CO.

LONDON:

TRÜBNER AND CO.

1860.

# CONTENTS.

JANUARY, 1860.

	PAGE		PAGE
PRIZE PROBLEMS FOR STUDENTS.....	105	ALSO OF THE FIFTEEN PROBLEMS IN THE TANGENCIES OF SPHERES. By Prof. George W. Coakley. ....	116
REPORT OF THE JUDGES UPON THE SOLU- TIONS OF THE PRIZE PROBLEMS IN NO. I., Vol. II.....	106	INSTANCES OF NEARLY COMMENSURABLE PERIODS IN THE SOLAR SYSTEM. By Prof. Daniel Kirkwood.....	126
ANOTHER SOLUTION OF PRIZE PROBLEM I. No. IX., Vol. I. By George Eastwood..	111	THE ELEMENTS OF QUATERNIONS. By W. P. G. Bardett.....	128
NOTE ON DECIMALS. By Samuel P. Bates.	112	NOTES ON PROBABILITIES. By Simon New- comb.....	134
REDUCTION OF FRACTIONS TO A COMMON DENOMINATOR.....	113	A SECOND BOOK IN GEOMETRY. By Rev. Thomas Hill.....	140
NOTE ON THE SUPERIOR LIMIT OF THE ROOTS OF AN EQUATION. By E. B. Weaver.....	114	EDITORIAL ITEMS.....	143
ANALYTICAL SOLUTIONS OF THE TEN PRO- BLEMS IN TANGENCIES OF CIRCLES; AND			

## ADVERTISEMENTS

Of interest to scientific and literary men, will be received *by the publishers*, up to the 20th of each month, for the next Number. Advertisers, to insure accuracy, should write distinctly, and on one side only, of paper.

### TERMS.

One Page, first insertion, . . . . .	\$ 15.00.
“ “ each subsequent insertion, . . . . .	12.00.
Half Page, first insertion, . . . . .	9.00.
“ “ each subsequent insertion, . . . . .	7.00.
Quarter Page, first insertion, . . . . .	5.00.
“ “ each subsequent insertion, . . . . .	4.00.

All Advertisements not otherwise marked are considered as ordered inserted till forbid.

Entered according to Act of Congress in the year 1860, by

J. D. RUNKLE,

In the Clerk's Office of the District Court of the District of Massachusetts.

### TERMS.

A single Copy, . . . . .	\$ 3.00 per annum, in advance.
“ “ . . . . .	4.00 “ after 6 months.
Two Copies to one Address, . . . . .	5.00 “ in advance.
Five Copies “ “ . . . . .	11.00 “ “
Ten Copies “ “ . . . . .	20.00 “ “

\* \* To any one getting up a Club of Ten Subscribers, and remitting the \$20.00, we will give a copy *gratis*. If Vol. I. is preferred, it will be given *gratis* by the Editor. Parties will please state distinctly which Volume they prefer.

IVISON, PHINNEY, & CO., *Publishers, New York.*

CAMBRIDGE: WELCH, BIGELOW, AND CO., ELECTROTYPERS AND PRINTERS.







THE

# MATHEMATICAL MONTHLY.

Vol. II . . . JANUARY, 1860. . . . No. IV.

PRIZE PROBLEMS FOR STUDENTS.

I. A and B can do a piece of work in  $m$  days; B and C in  $n$  days; in what time can A and C do the same, it being supposed that A can do  $p$  times as much as B in a given time?

II. Show that  $\frac{2+\sqrt{3}}{\sqrt{2}+\sqrt{2+\sqrt{3}}} + \frac{2-\sqrt{3}}{\sqrt{2}-\sqrt{2-\sqrt{3}}} = \sqrt{2}$ .

III. Find the roots of the equation  $x^3 - 6x = 4$ , by trigonometry.

IV. Four persons, A, B, C, D, in order, beginning with A, cut a pack of cards, replacing them after each cut, on condition that the first who cuts a heart shall win. What are their respective probabilities of success?

V. The notation of Problem V. in the November number of the Monthly being retained, prove that in the plane

$$\cot \frac{1}{2} A + \cot \frac{1}{2} B + \cot \frac{1}{2} C = \cot \frac{1}{2} A \cot \frac{1}{2} B \cot \frac{1}{2} C;$$

and in the sphere

$$\cot \frac{1}{2} A + \cot \frac{1}{2} B + \cot \frac{1}{2} C = \cot \frac{1}{2} A \cot \frac{1}{2} B \cot \frac{1}{2} C \frac{\cos(\rho + \delta) \cos(\rho - \delta)}{\cos^2 r \cos^2 \rho},$$

$$\cot \frac{1}{2} A \frac{\cos \delta'}{\cos \rho} + \cot \frac{1}{2} B \frac{\cos \delta''}{\cos \rho''} + \cot \frac{1}{2} C \frac{\cos \delta'''}{\cos \rho'''} = \cot \frac{1}{2} A \cot \frac{1}{2} B \cot \frac{1}{2} C \frac{\cos \delta}{\cos \rho}.$$

The solutions of these problems must be received by March 1, 1860.

REPORT OF THE JUDGES UPON THE SOLUTIONS OF THE  
PRIZE PROBLEMS IN No. I., Vol. II.

The first Prize is awarded to J. D. VAN BUREN, Rensselaer Polytechnic Institute, Troy, N. Y.

The second Prize is awarded to O. B. WHEELER, Sophomore class, University of Michigan, at Ann Arbor.

The third Prize is awarded to GEORGE H. TOWER, Classical High School, Petersham, Mass.

PRIZE SOLUTION OF PROBLEM II.

By PERRIN B. PAGE, Nunda, N. Y.

In any plane triangle, prove that the sines of the angles are inversely as the perpendiculars let fall from them upon the opposite sides.

Let  $ABC$  be the triangle, and denote by  $a, b, c$ , the perpendiculars dropped respectively from the angles  $A, B, C$ . By trigonometry

$$(1.) \quad \frac{\sin A}{BC} = \frac{\sin B}{AC} = \frac{\sin C}{AB}.$$

By geometry, twice the area of  $ABC$  is

$$(2.) \quad BC \times a = AC \times b = AB \times c.$$

Multiplying equations (1) and (2) together, member by member, gives,


$$\sin A \times a = \sin B \times b = \sin C \times c; \text{ or,}$$

$$\sin A : \sin B : \sin C = c : b : a.$$

SECOND SOLUTION. — By definition,

$$\sin A = \frac{b}{AB} = \frac{c}{AC}, \quad \sin B = \frac{a}{AB} = \frac{c}{BC},$$

$$\therefore \frac{\sin A}{\sin B} = \frac{b}{a} = \frac{BC}{AC}.$$



$$\therefore \sin A : \sin B = b : a = BC : AC,$$

which not only proves the proposition, but also that *the sines of the angles are proportional to their opposite sides*. This is a slight modification of W. C. HENCK'S solution.

#### PRIZE SOLUTION OF PROBLEM III.

By ISAAC H. TURRELL, Mt. Carmel, Indiana.

Having given the diagonals of a quadrilateral inscribed in a given circle to determine its sides geometrically, when the diagonals intersect each other at right angles. — Communicated by Professor D. W. HOYT.

Let  $AB$  and  $CD$  be the given diagonals. Inscribe  $AB$  in the given circle. Next, draw parallel to  $AB$  and at a distance from the centre equal to one half the other diagonal  $CD$ , the line  $CC'$ . From either extremity of this line, as  $C$ , draw a line perpendicular to  $AB$  and produce it till it meets the circumference in  $D$ . The line  $CD$  is therefore inscribed in the circle, perpendicular to  $AB$ , and  $ACBD$  is the required quadrilateral. CHARLES B. BOUTELLE'S solution is essentially the same as the above.



#### PRIZE SOLUTION OF PROBLEM IV.

By J. D. VAN BUREN, Rensselaer Polytechnic Inst., Troy, N. Y.

Given

$$\begin{aligned} (1.) \quad xw + yz &= ab, & (2.) \quad xy + zw &= ad \\ (3.) \quad xz + yw &= bd, & (4.) \quad x^2 + w^2 &= y^2 + z^2; \end{aligned}$$

to find the values of  $x, y, z$ , and  $w$ . — Communicated by Professor D. W. HOYT.

Adding and subtracting (1) and (3), (1) and (2), and also multiplying (3) and (2) together, we obtain

$$\begin{aligned} (5.) \quad x + y &= \frac{bd + ab}{z + w}, & (7.) \quad x + z &= \frac{ab + ad}{w + y}, \\ (6.) \quad x - y &= \frac{bd - ab}{z - w}, & (8.) \quad x - z &= \frac{ab - ad}{w - y}, \end{aligned}$$

$$(9.) \quad yz(x^2 + w^2) + xw(z^2 + y^2) = ab d^2.$$

Combining the products of (5) and (6), and (7) and (8), with (9) and (4) respectively, we get

$$x^2 - y^2 = \pm b \sqrt{d^2 - a^2}, y^2 - w^2 = \pm a \sqrt{b^2 - d^2}, x^2 + w^2 = d^2 = y^2 + z^2.$$

Therefore,

$$2x^2 = d^2 \pm b \sqrt{d^2 - a^2} \pm a \sqrt{b^2 - d^2},$$

$$2y^2 = d^2 \mp b \sqrt{d^2 - a^2} \pm a \sqrt{b^2 - d^2},$$

$$2w^2 = d^2 \mp b \sqrt{d^2 - a^2} \mp a \sqrt{b^2 - d^2},$$

$$2z^2 = d^2 \pm b \sqrt{d^2 - a^2} \mp a \sqrt{b^2 - d^2}.$$

Whence we obtain the values of the required quantities by dividing by 2 and extracting the square root. Similar solutions were given by WILLIAM W. JOHNSON and O. B. WHEELER.

O. B. WHEELER remarks that the equations of Problem IV. are readily obtained from the figure of Problem III. From geometry we have, "*The rectangle of the two diagonals is equivalent to the sum of the rectangles of the opposite sides,*" which is (1). We have (2) and (3) by the proposition, "*In any triangle the rectangle contained by two sides is equivalent to the rectangle contained by the diameter of the circumscribed circle and the perpendicular let fall on the third side;*" and (4) from the sum of the equations

$$AE^2 = w^2 - DE^2 = y^2 - EC^2,$$

$$BE^2 = x^2 - EC^2 = z^2 - DE^2.$$

The connection between Problems III. and IV. was also noticed by W. C. CLEVELAND, who gave the following geometrical demonstration of (4).

Let the chord  $CC'$  be drawn parallel to  $AB$ ; then  $AC' = BC = x$ , angle  $C'CD = 90^\circ$ ;

$$\therefore C'D \text{ is a diameter; } \therefore x^2 + w^2 = d^2.$$

Similarly  $y^2 + z^2 = d^2$ ; whence equation (4).

PRIZE SOLUTION OF PROBLEM V.

By J. D. VAN BUREN, Rensselaer Polytechnic Institute, Troy, N. Y.

If, in a plane or spherical triangle,  $A, B, C$  denote the angles, and  $a, b, c$  the opposite sides respectively; if  $r, \rho$  denote the radii of the circumscribed and inscribed circles, and  $\delta$  the distance between the centres of these circles; then in the plane triangle

$$\sin A + \sin B + \sin C = 4 \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C,$$

and in the spherical triangle

$$\sin A + \sin B + \sin C = 4 \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C \frac{\cos \delta}{\cos r \cos \rho}.$$

*Plane Triangle.* We have from plane trigonometry,

$$\sin A = \frac{2k}{bc}, \quad \sin B = \frac{2k}{ac}, \quad \sin C = \frac{2k}{ab},$$

when  $k = \sqrt{s(s-a)(s-b)(s-c)}$ , and  $s = \frac{1}{2}(a+b+c)$ .

$$\therefore \sin A + \sin B + \sin C = \frac{2k(a+b+c)}{abc} = \frac{4ks}{abc}.$$

$$\text{Also, } \cos \frac{1}{2} A = \sqrt{\frac{s(s-a)}{bc}}, \quad \cos \frac{1}{2} B = \sqrt{\frac{s(s-b)}{ac}}, \quad \cos \frac{1}{2} C = \sqrt{\frac{s(s-c)}{ab}}.$$

$$\therefore \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C = \frac{ks}{abc},$$

$$\therefore \sin A + \sin B + \sin C = 4 \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C.$$

*Spherical Triangle.* We have, from spherical trigonometry,

$$\sin A = \frac{2k}{\sin b \sin c}, \quad \sin B = \frac{2k}{\sin a \sin c}, \quad \sin C = \frac{2k}{\sin a \sin b},$$

in which

$$k = \sqrt{\sin s \sin(s-a) \sin(s-b) \sin(s-c)}, \text{ and } s = \frac{1}{2}(a+b+c).$$

$$(1.) \quad \therefore \sin A + \sin B + \sin C = \frac{2k(\sin a + \sin b + \sin c)}{\sin a \sin b \sin c}.$$

Also,

$$\cos \frac{1}{2} A = \sqrt{\frac{\sin s \sin(s-a)}{\sin b \sin c}}, \quad \cos \frac{1}{2} B = \sqrt{\frac{\sin s \sin(s-b)}{\sin a \sin c}},$$

$$\cos \frac{1}{2} C = \sqrt{\frac{\sin s \sin(s-c)}{\sin a \sin b}}.$$



$$\therefore \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C = \frac{k \sin s}{\sin a \sin b \sin c}.$$

But we have (CHAUVENET'S Trig., p. 251, and Eq. 319),

$$\frac{\cos \delta}{\cos r \sin \varphi} = \frac{\sin a + \sin b + \sin c}{2k}, \quad \tan \varphi = \frac{k}{\sin s},$$

$$\therefore \sin \varphi = \frac{k \cos \varphi}{\sin s} \quad \therefore \frac{\cos \delta}{\cos r \cos \varphi} = \frac{\sin a + \sin b + \sin c}{2 \sin s}.$$

$$(2.) \quad \therefore \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C \cdot \frac{\cos \delta}{\cos r \cos \varphi} = \frac{k (\sin a + \sin b + \sin c)}{2 \sin a \sin b \sin c}.$$

Therefore, by comparing (1) and (2) we get

$$\sin A + \sin B + \sin C = 4 \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C \cdot \frac{\cos \delta}{\cos r \cos \varphi}.$$

#### SECOND SOLUTION.

By O. B. WHEELER, University of Michigan.

*Plane Triangle.* Since in a plane triangle  $\frac{1}{2}(A + B + C) = \frac{1}{2}\pi$ , we have

$$(1.) \quad \cos \frac{1}{2}(A + B + C) = \cos \frac{1}{2}\pi = 0.$$

$$(2.) \quad \cos \frac{1}{2}(-A + B + C) = \cos(\frac{1}{2}\pi - A) = \sin A,$$

$$(3.) \quad \cos \frac{1}{2}(A - B + C) = \cos(\frac{1}{2}\pi - B) = \sin B,$$

$$(4.) \quad \cos \frac{1}{2}(A + B - C) = \cos(\frac{1}{2}\pi - C) = \sin C.$$

Taking the sum of (1) and (2), and also of (3) and (4), we obtain

$$(5.) \quad 2 \cos \frac{1}{2} A \cos \frac{1}{2}(B + C) = \sin A,$$

$$(6.) \quad 2 \cos \frac{1}{2} A \cos \frac{1}{2}(B - C) = \sin B + \sin C.$$

Adding (5) and (6), developing and reducing, we have

$$4 \cos \frac{1}{2} A \cos \frac{1}{2} B \cos \frac{1}{2} C = \sin A + \sin B + \sin C.$$

MR. WHEELER'S solution for the spherical triangle is essentially the same as the one already given.

SIMON NEWCOMB.

W. P. G. BARTLETT.

TRUMAN HENRY SAFFORD.

ANOTHER SOLUTION OF PRIZE PROBLEM I., No. IX.

By GEORGE EASTWOOD, Saxonville, Mass.

THERE are two ways of solving a geometrical problem *geometrically*. One way is to solve it by *synthesis*, the other way is to solve it by *analysis*. In solving a problem by analysis, we assume that the work is *done*; and then proceed, as the word implies, to take it to pieces, and to examine and develop the properties of each part and its relations to all the other parts. To solve the same problem by synthesis, we proceed to construct it upon elementary principles, from the given data and such combinations of them, as the exigencies of the case may require. We have a beautiful illustration of the synthetic method in Mr. EVANS's solution of Prize Problem I. of No. IX., published in the last Monthly. The following solution of the same problem is offered to young students as a simple specimen of the analytic method.\*



*Analysis.* Suppose the thing done, that  $ACB$  is the required triangle, and that  $AFBE$ ,  $GHI$ , are its circumscribing and inscribed circles. Join  $O, O'$ , the centres of the circles, and draw the radii  $O'G, O'H, O'I$ , to the points of contact  $G, H, I$ . Draw the diameter  $EF$  perpendicular to the hypotenuse  $AB$ , and  $CD$  perpendicular to  $EF$ .

By the question, we have given,

$$\begin{aligned} AC - BC &= (AG + GC) - (BH + HC) \\ &= AI - BI = 2OI = 2MN \text{ suppose.} \end{aligned}$$

---

\* Strictly speaking, the two ways of solving a geometrical problem, above indicated, are not two distinct and independent methods of doing the same thing, but rather the different parts of one full and perfect method. The one process is the reverse of the other, and the use of each is indispensable to a complete solution.

Also, we have given,

$$AB - BC = (AI + IB) - (IB + IO) = AI - IO,$$

which suppose equal to  $2LK$ . Take  $2MN$  from  $4LK$ , then

$$AI + BI - 2IO = 4LK - 2MN;$$

$$\therefore AO = 2LK - MN + IO.$$

By Prop. VI, *Liverpool Student*,  $EO \cdot FD = OI^2 = MN^2$ ; so that when  $EO = AO$  is found,  $FD$  will be given.

By Prop. 47, Euc. I.

$$\begin{aligned} OI^2 + OI^2 &= OO^2 = AO(AO - 2OI) \\ &= (2LK - MN)^2 - OI^2 = (2LK - OI)^2 - OI^2, \\ \therefore OI^2 &= 2LK(LK - MN). \end{aligned}$$

As  $LK$  and  $MN$  are both given lines, therefore  $OI$  is given. Hence this

*Construction.* Find (Euc. VI. 13)  $OI$  a mean proportional between  $2LK$  and  $LK - MN$ , and with centre  $O$  and radius  $OA = 2LK - MN + OI$ , describe a circle. Draw the horizontal and vertical diameters  $AB$ ,  $FE$ , and upon  $FE$  apply (Euc. VI. 11)  $FD$  a third proportional to  $EO$ ,  $OI$ . Draw  $DC$  parallel to  $AB$  to meet the circle in  $C$ , and join  $AC$ ,  $BC$ ; then  $ACB$  is the required triangle. The demonstration is evident from the analysis.

---

#### NOTES AND QUERIES.

1. *Note on Decimals.* — Decimals should be taught in written arithmetic in connection with whole numbers. Let them be treated in the same manner as whole numbers, and not as common fractions, there being no necessity for confusing the mind of the pupil by writing the denominator. In 31.3 there is no more need of indicating that the three standing in tenths' place is divided by ten, than

that the three standing in ten's place is multiplied by ten. The pupil can readily be made to understand these relations without the divisor or multiplier. There should be no separate divisions in arithmetic for Decimal Fractions and Federal Money. The four simple rules should embrace these, and Percentage and Interest should be treated immediately after as a development of the decimal notation. The tables and Compound Numbers should immediately precede Vulgar Fractions, — the former *increasing* in an irregular ratio, and the latter *decreasing* in an irregular ratio. If authors in writing arithmetics would observe this order, it would result in great advantage to the learner. — SAMUEL P. BATES, Superintendent of Public Instruction, Crawford County, Pa.

2. *Reduction of Fractions to a Common Denominator.* — After the pupil has thoroughly learned that multiplying or dividing both terms of a fraction by the same number does not change its value, he is then prepared to learn how to reduce fractions to a common denominator. The following process is very simple, and the pupil should be required to repeat it until it is thoroughly understood.

Let it be required, for example, to reduce  $\frac{7}{14}$ ,  $\frac{9}{15}$ ,  $\frac{23}{28}$ ,  $\frac{19}{35}$ ,  $\frac{13}{21}$  to a common denominator. First, resolve the terms of all the fractions

into their prime factors, and write them in a horizontal row. Now it is evident that when the denominators are common they will contain the same factors.

1st.	2d.	3d.	4th.	5th.
$\frac{7}{2 \cdot 7}$	$\frac{3 \cdot 3}{3 \cdot 5}$	$\frac{23}{2 \cdot 2 \cdot 7}$	$\frac{19}{5 \cdot 7}$	$\frac{2 \cdot 2 \cdot 3}{3 \cdot 7}$
$\frac{7}{2 \cdot 7}$	$\frac{3 \cdot 3 \cdot 7}{3 \cdot 5 \cdot 7}$	$\frac{23}{2 \cdot 2 \cdot 7}$	$\frac{19}{5 \cdot 7}$	$\frac{2 \cdot 2 \cdot 3}{3 \cdot 7}$
$\frac{7}{2 \cdot 7}$	$\frac{3 \cdot 7}{5 \cdot 7}$	$\frac{23}{2 \cdot 2 \cdot 7}$	$\frac{19}{5 \cdot 7}$	$\frac{2 \cdot 2}{7}$
$\frac{7 \cdot 5}{2 \cdot 7 \cdot 5}$	$\frac{3 \cdot 7}{5 \cdot 7}$	$\frac{23 \cdot 5}{2 \cdot 2 \cdot 7 \cdot 5}$	$\frac{19}{5 \cdot 7}$	$\frac{2 \cdot 2 \cdot 5}{7 \cdot 5}$
$\frac{2 \cdot 7 \cdot 5}{2 \cdot 2 \cdot 7 \cdot 5}$	$\frac{2 \cdot 2 \cdot 3 \cdot 7}{2 \cdot 2 \cdot 5 \cdot 7}$	$\frac{23 \cdot 5}{2 \cdot 2 \cdot 7 \cdot 5}$	$\frac{2 \cdot 2 \cdot 19}{2 \cdot 2 \cdot 5 \cdot 7}$	$\frac{2 \cdot 2 \cdot 2 \cdot 2 \cdot 5}{2 \cdot 2 \cdot 7 \cdot 5}$

Let us first compare them with reference to the factor 7. This factor is found in all the denominators but one; and in order

to make them common with reference to 7, it must either be introduced into the denominator of the 2d fraction, or removed from the denominators of all the other fractions. But to remove a factor from the denominator of a fraction without changing its value, it must be removed from the numerator also. Now 7 is a factor in only one of the numerators, and the denominators cannot be made common by removing the 7. It must therefore be introduced into the denominator of the second fraction to make the denominators common; and also into the numerator, in order not to change the value of the fraction. We thus get the second row. Next compare the denominators with reference to the factor 3. It is found in only two of them, from which it may be removed, since it is also found in the corresponding numerators. Thus we get the third row. (It is plain, that instead of removing the factor 3 from the 2d and 5th fractions, the denominators might be made common with reference to 3 by introducing it into the terms of all the other fractions. We should not, however, in this way reduce the fractions to their least common denominator.) Next introduce the factor 5 and get the fourth row; and lastly, introduce the factors 2. 2, which makes all the denominators common. — TEACHER.

3. *Note on the superior limit of the Roots of an Equation.*

PROP. I. *The greatest negative coefficient of an equation, plus unity, is a superior limit of its roots.*

PROOF. Let us assume the general equation,

$$x^n \pm Ax^{n-1} \pm Bx^{n-2} \dots \pm Nx^{n-r+1} \pm V = 0.$$

A superior limit of the roots of an equation must produce a positive result when substituted for  $x$ ; that is, the sum of all the positive terms must exceed the sum of all the negative ones by some quantity  $R$ . The most unfavorable case evidently is that in which *all the terms after the first are negative and have equal coefficients*. Under this assumption our equation becomes



$$x^n - Nx^{n-1} - Nx^{n-2} \dots - N = 0.$$

Suppose  $a$  to be greater than the greatest root, then

$$a^n - Na^{n-1} - Na^{n-2} \dots - N = R; \text{ or}$$

$$a^n - N(a^{n-1} - a^{n-2} \dots - 1) = R; \text{ or}$$

$$a^n - N\left(\frac{a^n - 1}{a - 1}\right) = R.$$

$$\therefore a^n = N\left(\frac{a^n - 1}{a - 1}\right) + R = \frac{Na^n}{a - 1} - \frac{N}{a - 1} + R.$$

Now suppose  $a$  to have been so taken that  $\frac{N}{a - 1} = R$ , as may always be done; then we shall have

$$a^n = \frac{Na^n}{a - 1}, \quad \text{or } a - 1 = N, \quad \text{or } a = N + 1.$$

But  $a > x$ ,  $\therefore N + 1 > x$ ; and hence  $N + 1$  is a superior limit, as was to be proved.

PROP. 2. *If unity be added to that root of the greatest negative coefficient, which is denoted by the number of terms preceding the first negative term, the result will be a superior limit of the roots of the equation.*

In the equation

$$x^n + Ax^{n-1} + Bx^{n-2} + \dots - Kx^{n-r} \pm \dots - Px^{n-r-s} \dots \pm V = 0,$$

suppose  $K$  to be the *first*, and  $P$  the *greatest*, negative coefficient, and  $r$  the number of terms which precede the first negative term. Evidently the most unfavorable case is that in which the coefficients  $A, B, C$ , &c., preceding  $K$  are zero, and all the coefficients  $K, L, M$ , &c., are all negative and equal. Under this supposition our equation becomes

$$x^n - Px^{n-r} - Px^{n-r-1} \dots - P = 0.$$

Suppose  $a$  to be greater than the greatest root, then

$$a^n - Pa^{n-r} - Pa^{n-r-1} \dots - P = R; \text{ or}$$

$$a^n - P\left(\frac{a^{n-r+1} - 1}{a - 1}\right) = R,$$

$$\therefore a^n = P \left( \frac{a^{n-r+1}-1}{a-1} \right) + R = \frac{Pa^{n-r+1}}{a-1} - \frac{P}{a-1} + R.$$

Suppose  $a$  to have been so taken that  $\frac{P}{a-1} = R$ , as may always be done; then we shall have

$$a^n = \frac{Pa^{n-r+1}}{a-1}, \text{ or } (a-1)a^{n-r+1} = P. \text{ But } a-1 < a;$$

$$\therefore (a-1)^{r-1} < a^{r-1}, (a-1)^r < (a-1)a^{r-1}.$$

Therefore, since  $(a-1)a^{r-1} = P$ ,  $(a-1)^r < P$ ;

$$\therefore a-1 < \sqrt[r]{P}, \text{ or } a < \sqrt[r]{P} + 1.$$

But  $a > x$ ,  $\therefore \sqrt[r]{P} + 1 > x$ , as was to be proved. — E. B. WEAVER, Normal School, Lancaster Co., Pa.

# ANALYTICAL SOLUTIONS "OF THE TEN PROBLEMS IN THE TANGENCIES OF CIRCLES; AND ALSO OF THE FIFTEEN PROBLEMS IN THE TANGENCIES OF SPHERES."

By GEORGE W. COAKLEY, Professor of Mathematics, College of St. James, Washington Co., Md.

PART of the title of this article is derived from that of Major ALVORD's paper in the eighth volume of the "*Smithsonian Contributions to Knowledge*." The analytic solutions here given were obtained in the winter of 1856, soon after I became acquainted with Major ALVORD's paper; they were attempted in consequence of the statements in that paper, that no mathematician had yet succeeded in reducing the analytical solution of these problems to equations of the second degree. At the end of his geometrical solutions, Major ALVORD states, that the algebraic solution leads to an equation of the *eighth degree* when applied to the most general problem of the tangencies of circles; and to an equation of the *sixteenth degree* when applied to the most general problem of the tangencies of spheres. It will be found, however, that in this article both of these general

problems, which include all the others contained in the Smithsonian paper, are solved completely by equations not surpassing the second degree. The solution of the *particular* problems included in the two *general* problems here treated may be obtained either by applying to them the general equations developed in this article; or by treating them separately in a similar manner, the process being of still easier application to most of these particular problems. Thus the whole twenty-five problems of the Smithsonian paper may be considered as reduced to equations of the second degree, when the two most comprehensive problems have been so reduced.

#### FIRST GENERAL PROBLEM.

*"To draw a Circle tangent to three given Circles."*

Let the radii of the given circles be denoted by  $r_1, r_2, r_3$ , respectively, and let the corresponding co-ordinates of their centres be denoted by  $(a_1, b_1), (a_2, b_2), (a_3, b_3)$ ; also let  $\rho, (\alpha, \beta)$  be the radius and centre of the required circle, all being referred to any system of rectangular co-ordinate axes in the common plane of the circles. The equations of these circles will be respectively

$$(1.) \quad (x - a_1)^2 + (y - b_1)^2 = r_1^2,$$

$$(2.) \quad (x - a_2)^2 + (y - b_2)^2 = r_2^2,$$

$$(3.) \quad (x - a_3)^2 + (y - b_3)^2 = r_3^2,$$

$$(4.) \quad (x - \alpha)^2 + (y - \beta)^2 = \rho^2,$$

Let  $x_1, y_1$  be the point of contact of (1) and (4),

"  $x_2, y_2$  " " " (2) and (4),

"  $x_3, y_3$  " " " (3) and (4).

Then the point  $x_1, y_1$  must satisfy the conditions,

$$(5.) \quad (x_1 - a_1)^2 + (y_1 - b_1)^2 = r_1^2,$$

$$(6.) \quad (x_1 - \alpha)^2 + (y_1 - \beta)^2 = \rho^2,$$

$$(7.) \quad \frac{dy_1}{dx_1} = -\frac{x_1 - a_1}{y_1 - b_1} = -\frac{x_1 - \alpha}{y_1 - \beta},$$

and similar conditions exist for the circles (2) and (3) combined with (4); but they may readily be derived from (5), (6), and (7) by simply changing the subscript 1 into 2 or 3.

From (7) we have

$$\frac{(x_1 - a_1)^2}{(y_1 - b_1)^2} + 1 = \frac{(x_1 - \alpha)^2}{(y_1 - \beta)^2} + 1,$$

$$\therefore \frac{r_1^2}{(y_1 - b_1)^2} = \frac{\rho^2}{(y_1 - \beta)^2}.$$

$$(8.) \quad \therefore y_1 - \beta = \pm \frac{\rho}{r_1} (y_1 - b_1),$$

$$(9.) \quad x_1 - \alpha = \pm \frac{\rho}{r_1} (x_1 - a_1),$$

$$\therefore y_1 = \frac{r_1 \beta \mp b_1 \rho}{r_1 \mp \rho}, \quad x_1 = \frac{r_1 \alpha \mp a_1 \rho}{r_1 \mp \rho},$$

$$\therefore y_1 - b_1 = \frac{r_1 (\beta - b_1)}{r_1 \mp \rho}, \quad x_1 - a_1 = \frac{r_1 (\alpha - a_1)}{r_1 \mp \rho}.$$

Hence equation (5) becomes

$$(\alpha - a_1)^2 + (\beta - b_1)^2 = (r_1 \mp \rho)^2,$$

$$(A.) \quad \text{and in like manner } (\alpha - a_2)^2 + (\beta - b_2)^2 = (r_2 \mp \rho)^2,$$

$$(\alpha - a_3)^2 + (\beta - b_3)^2 = (r_3 \mp \rho)^2.$$

These equations evidently express the fact that the distance between the centres of any two of the tangent circles is equal to the sum or difference of their radii according as the contact is external or internal. It is also evident that the following are all the cases that can possibly occur, viz.:—

The required circle touches the circles (1), (2), (3) externally,			
"	"	"	" (1), (2), (3) internally,
"	"	"	" (1), (2), ext. and (3) int.,
"	"	"	" (1), (3), ext. and (2) int.,

The required circle touches the circles (2), (3), ext. and (1) int.,

“ “ “ “ (1), (2), int. and (3) ext.,

“ “ “ “ (1), (3), int. and (2) ext.,

“ “ “ “ (2), (3), int. and (1) ext.,

making eight cases, (the number found by Major ALVORD,) each of which is solved by a proper combination of the algebraic signs in the equations (A). In order to embrace all the cases in one solution, let  $R_1 = \pm r_1$ ,  $R_2 = \pm r_2$ ,  $R_3 = \pm r_3$ ; then equations (A) may be written

$$\begin{aligned} & (\alpha - a_1)^2 + (\beta - b_1)^2 = \{\pm (\rho - R_1)\}^2 \\ (A'.) \quad & (\alpha - a_2)^2 + (\beta - b_2)^2 = \{\pm (\rho - R_2)\}^2 \\ & (\alpha - a_3)^2 + (\beta - b_3)^2 = \{\pm (\rho - R_3)\}^2. \end{aligned}$$

To find the values of the unknown quantities,  $\alpha$ ,  $\beta$ ,  $\rho$ , let

$$(10.) \quad \alpha = A_1 \rho + A_2, \quad \beta = B_1 \rho + B_2,$$

$A_1$ ,  $A_2$ ,  $B_1$ ,  $B_2$  being functions of the *given* quantities, readily determined in each case by equations of the *first degree* in the following manner.

Developing the squares in the three equations (A'), and subtracting in turn the members of the first from those of the second and third, we have

$$(11.) \quad \begin{aligned} 2(a_1 - a_2)\alpha + 2(b_1 - b_2)\beta &= 2(R_1 - R_2)\rho + a_1^2 - a_2^2 + b_1^2 - b_2^2 - (R_1^2 - R_2^2), \\ 2(a_1 - a_3)\alpha + 2(b_1 - b_3)\beta &= 2(R_1 - R_3)\rho + a_1^2 - a_3^2 + b_1^2 - b_3^2 - (R_1^2 - R_3^2). \end{aligned}$$

Substituting from (10), the values of  $\alpha$  and  $\beta$ , equations (11) become

$$\begin{aligned} 2\{(a_1 - a_2)A_1 + (b_1 - b_2)B_1\}\rho + 2(a_1 - a_2)A_2 + 2(b_1 - b_2)B_2 &= \\ 2(R_1 - R_2)\rho + a_1^2 - a_2^2 + b_1^2 - b_2^2 - (R_1^2 - R_2^2), \\ 2\{(a_1 - a_3)A_1 + (b_1 - b_3)B_1\}\rho + 2(a_1 - a_3)A_2 + 2(b_1 - b_3)B_2 &= \\ 2(R_1 - R_3)\rho + a_1^2 - a_3^2 + b_1^2 - b_3^2 - (R_1^2 - R_3^2). \end{aligned}$$

Now the two equations (11) are insufficient to determine the values of the three unknown quantities,  $\alpha$ ,  $\beta$ ,  $\rho$ ; hence the result of the



preceding substitutions must be to give *identical equations*, or those capable of being verified by any value of  $\rho$  whatever; hence the coefficients of the corresponding powers of  $\rho$  must be equal.

Therefore

$$(12.) \quad \begin{aligned} (a_1 - a_2) A_1 + (b_1 - b_2) B_1 &= R_1 - R_2 \\ (a_1 - a_3) A_1 + (b_1 - b_3) B_1 &= R_1 - R_3, \end{aligned}$$

$$(13.) \quad \begin{aligned} (a_1 - a_2) A_2 + (b_1 - b_2) B_2 &= \frac{1}{2} \{a_1^2 - a_2^2 + b_1^2 - b_2^2 - (R_1^2 - R_2^2)\}, \\ (a_1 - a_3) A_2 + (b_1 - b_3) B_2 &= \frac{1}{2} \{a_1^2 - a_3^2 + b_1^2 - b_3^2 - (R_1^2 - R_3^2)\}. \end{aligned}$$

Hence

$$(14.) \quad \begin{aligned} A_1 &= \frac{(R_1 - R_2)(b_1 - b_3) - (R_1 - R_3)(b_1 - b_2)}{(a_1 - a_2)(b_1 - b_3) - (a_1 - a_3)(b_1 - b_2)} \\ B_1 &= \frac{(R_1 - R_3)(a_1 - a_2) - R_1 - R_2)(a_1 - a_3)}{(a_1 - a_2)(b_1 - b_3) - (a_1 - a_3)(b_1 - b_2)}, \end{aligned}$$

and by putting

$$\begin{aligned} D &= a_1^2 - a_2^2 + b_1^2 - b_2^2 - (R_1^2 - R_2^2) \\ E &= a_1^2 - a_3^2 + b_1^2 - b_3^2 - (R_1^2 - R_3^2) \\ F &= (a_1 - a_2)(b_1 - b_3) - (a_1 - a_3)(b_1 - b_2), \end{aligned}$$

we also get

$$(15.) \quad A_2 = \frac{(b_1 - b_3) D - (b_1 - b_2) E}{2 F}, \quad B_2 = \frac{(a_1 - a_2) E - (a_1 - a_3) D}{2 F}.$$

Thus the coefficients,  $A_1, B_1, A_2, B_2$ , of equations (10) are entirely determined in terms of the given quantities. From equations (10) we have

$$(16.) \quad \begin{aligned} \alpha - a_1 &= A_1(\rho - R_1) + A_2 + A_1 R_1 - a_1 = A_1(\rho - R_1) + m \\ \beta - b_1 &= B_1(\rho - R_1) + B_2 + B_1 R_1 - b_1 = B_1(\rho - R_1) + n, \end{aligned}$$

by making

$$(17.) \quad m = A_2 + A_1 R_1 - a_1, \quad n = B_2 + B_1 R_1 - b_1.$$

Hence the first equation of (A') becomes

$$(18.) \quad (\rho - R_1)^2 - \frac{2(m A_1 + n B_1)}{1 - (A_1^2 + B_1^2)} (\rho - R_1) = \frac{m^2 + n^2}{1 - (A_1^2 + B_1^2)}.$$

Hence

$$(19.) \quad \rho = R_1 + \frac{m A_1 + n B_1 \pm \sqrt{m^2 + n^2 - (n A_1 - m B_1)^2}}{1 - (A_1^2 + B_1^2)}.$$

Thus the equations (10), (14), (15), (17), and (19) completely solve the problem for every possible case.

It is evident that the problem is impossible whenever

$$(n A_1 - m B_1)^2 > m^2 + n^2;$$

and that of the two roots in (19), the one which gives a negative value for  $\rho$  is to be interpreted as giving a circle of contact the reverse of that implied by the values assumed for  $R_1$ ,  $R_2$ , and  $R_3$ . If, for example, the value assumed for  $R_2$  implied *external* contact, then the negative value of  $\rho$  would indicate *internal* contact with the second given circle.

I shall apply the preceding formulæ to determine the circle which passes through three given points, regarded as circles of infinitely small radius. Let the axis of abscissas pass through two of the given points,  $(a_2, b_2)$  and  $(a_3, b_3)$ , and let the origin bisect the distance between them; hence  $a_3 = -a_2$ ,  $b_2 = b_3 = 0$ ; also,  $r_1 = r_2 = r_3 = 0 = R_1 = R_2 = R_3$ .

Hence  $A_1 = 0, \quad B_1 = 0, \quad A_2 = 0,$

$$B_2 = \frac{1}{2} \frac{a_1^2 + b_1^2 - a_2^2}{b_1}; \quad m = -a_1, \quad n = \frac{a_1^2 - b_1^2 - a_2^2}{2 b_1},$$

$$\rho = \pm \sqrt{m^2 + n^2} = \pm \sqrt{\frac{4 a_1^2 b_1^2 + (a_1^2 - b_1^2 - a_2^2)^2}{4 b_1^2}};$$

$$\alpha = 0, \beta = \frac{1}{2} \frac{a_1^2 - b_1^2 - a_2^2}{b_1}.$$

The remaining point,  $(a_1, b_1)$  is thus far subject to no restriction; it is anywhere on the plane of the co-ordinate axes. If now we suppose first that it is on the same straight line with the other two points, then  $b_1 = 0$ ; hence  $\beta = \infty$ ,  $\rho = \pm \infty$ . Hence the radius

of the circle passing through three given points on the same straight line is infinite, and its centre is at an infinite distance from that line. The double sign of  $\rho$  shows that the contact may be considered in this case either external or internal. Secondly, if we suppose the point  $(a_1, b_1)$  to be on the axis of ordinates, at a distance from the origin equal to  $a_2$ , then  $a_1 = 0$ ,  $b_1 = a_2$ ,  $\beta = 0$ ,  $\alpha = 0$ ,  $m = 0$ ,  $n = -a_2$ ,  $\rho = \pm a_2$ , the circle having either external or internal contact with the three given points; and it is obvious that the results are correct.

As a second example, let it be required to find the circle which shall touch *externally* the three circles whose radii and centres are as follows:

$$\begin{array}{llll} r_1 = 1, & r_2 = 3, & r_3 = 2, & \text{hence } R_1 = -r_1 = -1, \\ a_1 = 0, & a_2 = 5, & a_3 = -3, & \text{" } R_2 = -r_2 = -3, \\ b_1 = 0, & b_2 = 0, & b_3 = 4, & \text{" } R_3 = -r_3 = -2. \end{array}$$

It will be found that  $A_1 = -0.4$ ,  $B_1 = -0.55$ ,  $A_2 = 1.7$ ,  $B_2 = 4.025$ ,  $m = 2.1$ ,  $n = 4.575$ ,  $\rho = 2.03672$ ,  $\alpha = 0.885312$ ,  $\beta = 2.904804$ , and the equations (A), or (A'), will be found to be satisfied.

#### SECOND GENERAL PROBLEM.

*"To draw a Sphere tangent to four given Spheres."*

Let the radii of the given spheres be denoted by  $r_1, r_2, r_3, r_4$ , and the co-ordinates of their centres by  $(a_1, b_1, c_1)$ ,  $(a_2, b_2, c_2)$ ,  $(a_3, b_3, c_3)$ ,  $(a_4, b_4, c_4)$ ; let  $\rho$  ( $\alpha, \beta, \gamma$ ) be the radius and centre of the required sphere; all being referred to rectangular co-ordinate axes in space.

It is evident that by an analysis similar to that for the circles, the distance of the centre  $(\alpha, \beta, \gamma)$  from each of the centres  $(a_1, b_1, c_1)$ , &c., would be found equal to the sum or difference of the corresponding radii, according as the spheres had external or internal contact; hence to determine the four unknown quantities,  $\alpha, \beta, \gamma, \rho$ , we should have the four equations

$$\begin{aligned}
 & (\alpha - a_1)^2 + (\beta - b_1)^2 + (\gamma - c_1)^2 = (\varrho \pm r_1)^2, \\
 & (\alpha - a_2)^2 + (\beta - b_2)^2 + (\gamma - c_2)^2 = (\varrho \pm r_2)^2, \\
 \text{(C.)} \quad & (\alpha - a_3)^2 + (\beta - b_3)^2 + (\gamma - c_3)^2 = (\varrho \pm r_3)^2, \\
 & (\alpha - a_4)^2 + (\beta - b_4)^2 + (\gamma - c_4)^2 = (\varrho \pm r_4)^2.
 \end{aligned}$$

Or if, to embrace all the cases in one solution, we put  $R_1 = \pm r_1$ ,  $R_2 = \pm r_2$ , &c., the preceding equations may be written

$$\begin{aligned}
 & (\alpha - a_1)^2 + (\beta - b_1)^2 + (\gamma - c_1)^2 = (\varrho - R_1)^2, \\
 & (\alpha - a_2)^2 + (\beta - b_2)^2 + (\gamma - c_2)^2 = (\varrho - R_2)^2, \\
 \text{(C'.)} \quad & (\alpha - a_3)^2 + (\beta - b_3)^2 + (\gamma - c_3)^2 = (\varrho - R_3)^2, \\
 & (\alpha - a_4)^2 + (\beta - b_4)^2 + (\gamma - c_4)^2 = (\varrho - R_4)^2.
 \end{aligned}$$

Let

$$\text{(1.)} \quad \alpha = A_1 \varrho + A_2, \quad \beta = B_1 \varrho + B_2, \quad \gamma = C_1 \varrho + C_2.$$

Then, by an analysis precisely similar to that employed in the general problem of the contact of circles, the following equations will be found for determining the values of  $A_1, B_1, C_1, A_2, B_2, C_2$ .

For the sake of abbreviation, let

$$\begin{aligned}
 M_2 &= a_1^2 - a_2^2 + b_1^2 - b_2^2 + c_1^2 - c_2^2 - (R_1^2 - R_2^2), \\
 \text{(2.)} \quad M_3 &= a_1^2 - a_3^2 + b_1^2 - b_3^2 + c_1^2 - c_3^2 - (R_1^2 - R_3^2), \\
 M_4 &= a_1^2 - a_4^2 + b_1^2 - b_4^2 + c_1^2 - c_4^2 - (R_1^2 - R_4^2);
 \end{aligned}$$

then

$$\begin{aligned}
 & (a_1 - a_2) A_1 + (b_1 - b_2) B_1 + (c_1 - c_2) C_1 = R_1 - R_2, \\
 \text{(3.)} \quad & (a_1 - a_3) A_1 + (b_1 - b_3) B_1 + (c_1 - c_3) C_1 = R_1 - R_3, \\
 & (a_1 - a_4) A_1 + (b_1 - b_4) B_1 + (c_1 - c_4) C_1 = R_1 - R_4.
 \end{aligned}$$

$$\begin{aligned}
 & (a_1 - a_2) A_2 + (b_1 - b_2) B_2 + (c_1 - c_2) C_2 = \frac{1}{2} M_2, \\
 \text{(4.)} \quad & (a_1 - a_3) A_2 + (b_1 - b_3) B_2 + (c_1 - c_3) C_2 = \frac{1}{2} M_3, \\
 & (a_1 - a_4) A_2 + (b_1 - b_4) B_2 + (c_1 - c_4) C_2 = \frac{1}{2} M_4.
 \end{aligned}$$

In order to abridge the expressions for  $A_1, A_2$ , &c., resulting from these equations, let

$$(5.) \quad \begin{aligned} P_a &= \frac{(b_1 - b_3)(c_1 - c_4) - (b_1 - b_4)(c_1 - c_3)}{(b_1 - b_2)(c_1 - c_3) - (b_1 - b_3)(c_1 - c_2)}, \\ Q_a &= \frac{(b_1 - b_4)(c_1 - c_2) - (b_1 - b_3)(c_1 - c_4)}{(b_1 - b_2)(c_1 - c_3) - (b_1 - b_3)(c_1 - c_2)}. \end{aligned}$$

Changing  $b$  into  $a$ , and  $a$  into  $b$ , in these expressions we have

$$(6.) \quad \begin{aligned} P_b &= \frac{(a_1 - a_3)(c_1 - c_4) - (a_1 - a_4)(c_1 - c_3)}{(a_1 - a_2)(c_1 - c_3) - (a_1 - a_3)(c_1 - c_2)}, \\ Q_b &= \frac{(a_1 - a_4)(c_1 - c_2) - (a_1 - a_3)(c_1 - c_4)}{(a_1 - a_2)(c_1 - c_3) - (a_1 - a_3)(c_1 - c_2)}. \end{aligned}$$

Changing again  $c$  into  $b$ , and  $b$  into  $c$ , in the last formulæ, we have

$$(7.) \quad \begin{aligned} P_c &= \frac{(a_1 - a_3)(b_1 - b_4) - (a_1 - a_4)(b_1 - b_3)}{(a_1 - a_2)(b_1 - b_3) - (a_1 - a_3)(b_1 - b_2)}, \\ Q_c &= \frac{(a_1 - a_4)(b_1 - b_2) - (a_1 - a_3)(b_1 - b_4)}{(a_1 - a_2)(b_1 - b_3) - (a_1 - a_3)(b_1 - b_2)}. \end{aligned}$$

$$(8.) \quad \begin{aligned} \therefore A_1 &= \frac{(R_1 - R_2) P_a + (R_1 - R_3) Q_a + R_1 - R_4}{(a_1 - a_2) P_a + (a_1 - a_3) Q_a + a_1 - a_4}, \\ B_1 &= \frac{(R_1 - R_2) P_b + (R_1 - R_3) Q_b + R_1 - R_4}{(b_1 - b_2) P_b + (b_1 - b_3) Q_b + b_1 - b_4}, \\ C_1 &= \frac{(R_1 - R_2) P_c + (R_1 - R_3) Q_c + R_1 - R_4}{(c_1 - c_2) P_c + (c_1 - c_3) Q_c + c_1 - c_4}. \end{aligned}$$

$$(9.) \quad \begin{aligned} A_2 &= \frac{1}{2} \frac{M_2 P_a + M_3 Q_a + M_4}{(a_1 - a_2) P_a + (a_1 - a_3) Q_a + a_1 - a_4}, \\ B_2 &= \frac{1}{2} \frac{M_2 P_b + M_3 Q_b + M_4}{(b_1 - b_2) P_b + (b_1 - b_3) Q_b + b_1 - b_4}, \\ C_2 &= \frac{1}{2} \frac{M_2 P_c + M_3 Q_c + M_4}{(c_1 - c_2) P_c + (c_1 - c_3) Q_c + c_1 - c_4}. \end{aligned}$$

Equations (1) may be written as follows :

$$(10.) \quad \begin{aligned} \alpha - a_1 &= A_1 (\varrho - R_1) + k, \\ \beta - b_1 &= B_1 (\varrho - R_1) + m, \\ \gamma - c_1 &= C_1 (\varrho - R_1) + n, \end{aligned}$$

by making



$$(11.) \quad \begin{aligned} k &= A_1 R_1 + A_2 - a_1, \\ m &= B_1 R_1 + B_2 - b_1, \\ n &= C_1 R_1 + C_2 - c_1. \end{aligned}$$

Then the first of equations (C') becomes

$$(\varrho - R_1)^2 - \frac{2(k A_1 + m B_1 + n C_1)}{1 - (A_1^2 + B_1^2 + C_1^2)} (\varrho - R_1) = \frac{k^2 + m^2 + n^2}{1 - (A_1^2 + B_1^2 + C_1^2)}.$$

Hence, by putting

$$G = (m A_1 - k B_1)^2 + (n A_1 - k C_1)^2 + (n B_1 - m C_1)^2,$$

we get

$$(12.) \quad \varrho = R_1 + \frac{k A_1 + m B_1 + n C_1 \pm \sqrt{k^2 + m^2 + n^2 - G}}{1 - (A_1^2 + B_1^2 + C_1^2)}.$$

Thus the problem is solved analytically by means of equations not transcending the *second degree*.

The number of cases of a sphere touching four given spheres may be determined as follows:

The required sphere may touch the four given spheres all externally, 1 case; or all internally, 1 case; or two internally, two externally, 6 cases; or three externally, one internally, 4 cases; or three internally, one externally, 4 cases; making sixteen cases, as found by Major ALVORD, all included in equations (C').

I must not omit to notice what would otherwise seem to be a defect in the preceding solutions. In both problems there seem to be exceptional cases to which the formulæ are inapplicable. For example, in the tangencies of circles, if  $b_1 = b_2 = b_3$ , then it appears that  $A_1 = \frac{0}{0}$ ,  $A_2 = \frac{0}{0}$ , and it is impossible to determine  $\varrho$ ,  $\alpha$ ,  $\beta$ , by the proposed method. The reason of this is obvious; for the assumed equations (10), in the first problem, were only employed because the *two equations* (11) contained *three* unknown quantities,  $\alpha$ ,  $\beta$ ,  $\varrho$ , and were, therefore, *indeterminate*. But when  $b_1 = b_2 = b_3$ ,

one of the unknown quantities,  $\beta$ , disappears from the equations (11); hence the assumed equations (10) are inapplicable, and, at the same time, they are not wanted; for  $\alpha$  and  $\rho$  are then determined by the two equations (11), each of the *first degree*, and then  $\beta$  is found from either of the equations (A') of the *second degree*. The same considerations apply to the corresponding case of the second general problem, when, for example,  $c_1 = c_2 = c_3 = c_4$ , in which case  $P_a = \frac{0}{0}$ ,  $Q_a = \frac{0}{0}$ ,  $P_b = \frac{0}{0}$ ,  $Q_b = \frac{0}{0}$ ; hence  $A_1$ ,  $B_1$ ,  $A_2$ , and  $B_2$  are indeterminate. But under these conditions the unknown quantity  $\gamma$  disappears from the three equations of the *first degree* obtained in the early part of the process; hence the remaining quantities,  $\alpha$ ,  $\beta$ ,  $\rho$ , are easily found, and thence  $\gamma$  from one of the equations (C'), of the *second degree*. Thus in every case the problems are brought within the resolution of equations not surpassing the second degree. I might now give various applications of the second problem which I have computed, but omit them for fear of occupying too much space in the Monthly unnecessarily.

---

#### INSTANCES OF NEARLY COMMENSURABLE PERIODS IN THE SOLAR SYSTEM.

By DANIEL KIRKWOOD, Professor of Mathematics, Indiana University.

THE following instances of nearly commensurable periods in the solar system have not, I believe, been previously noticed:—

For the 3d and 4th satellites of Jupiter we have

$$7 P^{\text{III}} - 3 P^{\text{IV}} = 22 \text{ m } 24 \text{ s} = \frac{1}{1071} P^{\text{IV}}; \dots (a).$$

For the Uranian System,

$$13 P^{\text{II}} - 4 P^{\text{IV}} = 26 \text{ m } 43 \text{ s} = \frac{1}{726} P^{\text{IV}}; \dots (b).$$

$$10 P^{\text{III}} - 21 P^{\text{II}} = 51 \text{ m } 6 \text{ s} = \frac{1}{245} P^{\text{III}}; \dots (c).$$

For the asteroids and major planets,

11 periods of Mars	=	5 periods of Astræa.
1 period of Jupiter	=	3 " Calliope.
3 periods "	=	10 " Clio.
4 " "	=	11 " Virginia.
6 " "	=	13 " Doris.
8 " "	=	29 " Ariadne.
9 " "	=	22 " Polyhymnia.
1 period of Saturn	=	9 " Ariadne.
2 periods "	=	13 " Leda.

We have also,

8 periods of Jupiter	=	7 periods of TUTTLE'S Comet.
6 " Saturn	=	13 " "
31 " "	=	12 " HALLEY'S Comet.
6 " Neptune	=	13 " "

*Remarks.*—The periods used in (a) are taken from HERSCHEL'S Outlines; those in (b) and (c) from the Proceedings of the American Association for the Advancement of Science, 1854, p. 55. The periods of the asteroids are taken from the Mathematical Monthly for February, 1859. The period of TUTTLE'S Comet is adopted from GOULD'S Astronomical Journal, No. 118. That of HALLEY'S is the mean of the six periods from 1378 to 1835. The epochs at which Neptune must have great perturbing influence on the motion of HALLEY'S Comet are separated by intervals of about 988 years. The last occurred in the early part of the fifteenth century;—the next will be about the beginning of the twenty-fifth. It may also be remarked that the planets Jupiter and Uranus present a case of almost exact commensurability. *Eighty-five times the period of the former, minus twelve times that of the latter, forms a quantity which amounts to only  $\frac{1}{1102}$  of the period of Uranus.* Hence in about  $1008\frac{1}{4}$  years from the epoch of any conjunction of these planets, another conjunction will occur only *twenty-three minutes* from the place of the former.

# THE ELEMENTS OF QUATERNIONS.

[Continued from Page 101.]

## IV. — SCALARS AND VECTORS.

19. If  $\angle q = \frac{\beta}{\alpha} = 0^\circ$ , or  $= 180^\circ$ , then  $\beta$  has either the same direction as  $\alpha$  or the opposite, and  $q$  degenerates into either a positive or negative real number, and is then called a *Scalar*. Like tensors, scalars may be applied to any line in space without regard to its direction; and like them, they are commutative in combination with any quaternion. The product of scalars is evidently a scalar; and the conjugate of a scalar does not differ from the scalar itself.

20. If  $\angle q = \pm 90^\circ$ , then  $q$  is called a *Vector*. Vectors will be denoted by  $v, v', \&c.$  It is evident from (4) and (3') that

$$(16) \quad K v = - v.$$

If  $T v = m$ , then by (11),

$$(17) \quad v^2 = - m^2;$$

that is, *the square of a vector is a negative scalar*. The square root of (17),  $v = m \sqrt{-1}$ , may be regarded as the expression for a vector of *indeterminate axis*, whose tensor is  $m$ . When  $m = 1$ ,  $v = \sqrt{-1}$  is called a *unit vector*.  $\sqrt{-1}_q$  may be used to denote this vector, when its axis is fixed in the direction of  $Ax.q$ . The geometrical signification then of  $\sqrt{-1}$  in this system, is the operation of turning any line to which it may be applied through an angle of  $90^\circ$  around any axis perpendicular to that line. Any quaternion may be expressed as a power of a vector, since, by (11), if  $m$  being a positive number,

$$(18) \quad q = [m \sqrt{-1}_q]^n, \quad \text{then } T q = m^n, \quad \angle q = n 90^\circ,$$

and the independent real quantities  $m$  and  $n$  are sufficient to determine  $T q$  and  $\angle q$ .

21. The lines  $\alpha$  and  $\beta$  being given, let the directions of the lines  $\gamma$  and  $\delta$  be respectively parallel and perpendicular to that of  $\alpha$ , and

let their lengths be such that  $\gamma + \delta = \beta$ . These conditions completely determine  $\gamma$  and  $\delta$ , both of which will be in the plane of  $\alpha$  and  $\beta$ . Then, by § 8,

$$q = \beta \div \alpha = (\gamma + \delta) \div \alpha = \gamma \div \alpha + \delta \div \alpha;$$

but  $\gamma \div \alpha$  is a scalar, and  $\delta \div \alpha$  is a vector. Any given quaternion then may be resolved into a sum of two other determinate quaternions, one of which is a scalar, and the other a vector; in this case the former is called the *scalar of the given quaternion*, and the latter *its vector*. The scalar of a quaternion,  $q$ , is written  $Sq$ , and its vector  $Vq$ ; thus

$$(19) \quad q = Sq + Vq = Vq + Sq.$$

From § 19, (16), and (4) we obtain

$$(20) \quad K Sq = Sq = SKq, \quad K Vq = -Vq = VKq; \text{ whence}$$

$$(19a) \quad Kq = Sq - Vq.$$

The sum and difference of (19) and (19a) gives

$$(21) \quad q + Kq = 2Sq, \quad \text{and} \quad q - Kq = 2Vq; \text{ or}$$

$$\text{symbolically,} \quad S = \frac{1}{2}(1 + K), \quad \text{and} \quad V = \frac{1}{2}(1 - K).$$

If two or more quaternions are equal, their scalars must be equal, and also their vectors; and conversely if their scalars and vectors are equal each to each, the quaternions must be equal.

22. Let the planes of the two vectors,  $v$  and  $v'$ , intersect in the line  $\alpha$ ; and let  $v\alpha = \beta$ , and  $v'\alpha = \beta'$ ; then both  $\beta$  and  $\beta'$  will be in the plane of  $Ax.v$  and  $Ax.v'$ , and, if  $\gamma = \beta + \beta'$ ,  $(v + v') = \gamma \div \alpha$ . But  $\gamma \div \alpha$  is a vector, whose axis, in the plane of  $Ax.v$  and  $Ax.v'$ , makes the same angles with these axes that the line  $\gamma$  does with the lines  $\beta$  and  $\beta'$  respectively, and whose tensor bears the same ratios to  $Tv$  and  $Tv'$  that  $T\gamma$  does to  $T\beta$  and  $T\beta'$  respectively. If then we set off on  $Ax.v$  and  $Ax.v'$  lines whose tensors are equal

respectively to  $Tv$  and  $Tv'$ , a parallelogram formed on these two sides will be similar to the parallelogram formed on  $\beta$  and  $\beta'$ , and its diagonal corresponding to the diagonal  $\gamma$  of the latter parallelogram, will be the axis of  $(v + v')$ , and the tensor of this diagonal will be equal to the tensor of  $(v + v')$ . Hereafter we shall consider *the axis of a vector as a line of definite length, whose tensor is equal to the tensor of the vector.* With this definition, then,

$$(22) \quad \Sigma Ax.v = Ax.\Sigma v.$$

23. Let the line  $\beta$  be the intersection of the planes of  $v'$  and  $v$ , and such that rotation around it is positive from  $Ax.v'$  to  $Ax.v$ ; and let  $v' = \alpha \div \beta$ , and  $v = \beta \div \gamma$ ; then both  $\alpha$  and  $\gamma$  will be in the plane of  $Ax.v'$  and  $Ax.v$ , and if we put

$$(23) \quad v'v = \alpha \div \gamma = q; \text{ then } Ax.q = \beta, \text{ and } \angle q = 180^\circ - \frac{Ax.v}{Ax.v'}.$$

Since  $Uv'v^{-1} = U(\alpha \div \beta)U(\beta \div -\gamma) = U(\alpha \div -\gamma)$ , if we put

$$v'v^{-1} = -Tv^{-2}.q = q'; \text{ then}$$

$$(23a) \quad Ax.q' = \beta, \quad \text{and } \angle q' = 180^\circ + \angle q = \frac{Ax.v'}{Ax.v}.$$

As a special case of (8), we have, by (16),

$$(24) \quad Kv'v = Kv.Kv' = vv'.$$

24. Let  $Ax.v$  be in the plane of  $q$ , and  $\alpha$  a line in the direction of  $Ax.q$ ; and let  $v\alpha = \beta$ , and  $q\beta = \gamma$ . Then

$$(25) \quad qv = (\gamma \div \beta)(\beta \div \alpha) = \gamma \div \alpha, \quad \text{a vector,}$$

$Tqv = Tq.Tv$ , and  $Ax.qv$  is so placed in the plane of  $q$  that  $\frac{Ax.qv}{Ax.v} = \frac{\gamma}{\beta} = \angle q$ . Therefore,

$$(26) \quad Ax.qv = qAx.v.$$

From (25), (16), and (8) we deduce

$$(27) \quad Kqv = -qv = Kv.Kq = -v.Kq, \quad qv = v.Kq;$$

and putting  $Kq$  for  $q$  in (27), we get, by (5),

$$(27a) \quad Kq \cdot v = v \cdot K^2 q = vq;$$

whence, making the same substitution in (26), we get by (27a)

$$(26a) \quad Ax.vq = Kq \cdot Ax.v.$$

Putting  $qv = v'$ , (26) becomes  $Ax.v' = qAx.v$ ; whence

$$(28) \quad q = v' \div v = v'v^{-1} = Ax.v' \div Ax.v, \text{ or, putting } v^{-1} \text{ for } v,$$

$$(28a) \quad v'v = Ax.v' \div Ax.v^{-1}.$$

25. Let the line  $\alpha^{-1}$ , the *reciprocal* of the line  $\alpha$ , be defined by the equations

$$(29) \quad \alpha = Ax.v, \quad \alpha^{-1} = Ax.v^{-1}; \text{ whence } (Ax.v)^{-1} = Ax.v^{-1}.$$

Equations (26) and (28a) give respectively by the substitution of (29)

$$(30) \quad Ax.qv^{-1} = q(Ax.v)^{-1},$$

$$(31) \quad v'v = Ax.v' \div (Ax.v)^{-1} = Ax.v' \cdot Ax.v.$$

26. The results expressed in (22), (28), (31), (26), and (30) show that a vector,  $v$ , is, under the definitions assumed, (in § 22 and (29),) perfectly represented in these cases by a line,  $Ax.v$ . Thus in § 22 the sum of vectors is proved *to be a vector*, and (22) expresses that the axis of this vector may be found by adding the axes of the given vectors. Equation (28) shows that the quotient of two vectors is the same quaternion as the quotient of their axes. Equation (31) shows that the definition (29) is so taken as to represent the multiplication of two vectors by the multiplication of their axes. Lastly, (25) shows that the product  $qv$  is, in that special case, *a vector*, and (26) shows that the axis of this vector may be found by multiplying  $Ax.v$  by  $q$ ; while (30) shows that the same property is true of  $qv^{-1}$ . The only possible cases remaining are the forms  $q + v$ ,  $qv$  and  $qv^{-1}$ , (except the special cases (25) and (30),)  $vq$  and  $v^{-1}q$ ;



and since these have, as yet, no meaning when a line, as  $Ax.v$ , is put in the place of  $v$ , we may now give them an arbitrary meaning in accordance with the conclusion, deduced from the other cases, that  $Ax.v$  may *always* be substituted for  $v$ ; that is, that a line is a vector, and that *whatever may be proved of lines is proved of the vectors of which they are the axes*. HAMILTON indeed introduces the term *vector* as a name for a straight line, deriving it from *vehere*, because a line is supposed to *carry* a point from one of its extremities to the other. Afterwards, assuming its other signification, as a special case of a quaternion, he proceeds somewhat in the reverse order to that in which the subject is developed here.

27. Any vector,  $v$ , may, by § 22, be expressed as the sum of three mutually rectangular vectors,  $v_x$ ,  $v_y$ , and  $v_z$ , whose axes are equal in length to the projections\* of  $Ax.v$  in their respective directions. If the tensors of  $v_x$ , &c., are denoted by  $(Tv)_x$ , &c., and their versors by  $i$ ,  $j$ , and  $k$  respectively, we have

$$q = Vq + Sq = (TVq)_x i + (TVq)_y j + (TVq)_z k + (Sq),$$

in which the four quantities in parentheses are four independent elements involved in the complete determination of  $q$ . Compare § 14.

28. The planes of the quaternions  $q = \beta \div \alpha$  and  $q' = \beta' \div \alpha$  intersect in the line  $\alpha$ ; and it is evident that  $Sq \cdot \alpha$  is the projection of  $\beta$  in the direction of  $\alpha$ , and  $Vq \cdot \alpha$  its projection on a plane perpendicular to  $\alpha$ ; whence it follows for the two quaternions,  $q$  and  $q'$ , and thence by easy extension for any number, that

$$(32) \quad S\Sigma = \Sigma S, \quad \text{and} \quad V\Sigma = \Sigma V.$$

But, as the value of  $SII$  does not depend on what or how many factors the product may be composed of, so long as the product

---

\* For greater completeness we may add that the *projection* of a line in any given direction, that is, on any given line, is so much of the latter line as is intercepted between perpendiculars dropped upon it from the extremities of the former.

itself remains the same; while  $II S$  evidently does depend on the composition of  $II$ ; it follows that we do *not* have in general  $S II = II S$ . And as § 23 shows that  $II V$  is not necessarily a vector, we do *not* have in general  $V II = II V$ .

29. We are now prepared to deduce the following relations, connecting the notation of quaternions with trigonometry. If  $q = \beta \div \alpha$ ,

$$(33) \quad S q \cdot T \alpha = T \beta \cdot \cos \alpha^\beta \quad S q = (T \beta \div T \alpha) \cos \alpha^\beta = T q \cdot \cos \alpha^\beta,$$

$$(33a) \quad V q \cdot T \alpha = T V q \cdot T \alpha \cdot U V q = T \beta \cdot \sin \alpha^\beta \cdot \sqrt{-1_q}, \quad V q = T q \cdot \sin \alpha^\beta \cdot \sqrt{-1_q};$$

whence

$$(34) \quad q = T q (\cos \alpha^\beta + \sin \alpha^\beta \cdot \sqrt{-1_q}), \quad U q = \cos \alpha^\beta + \sin \alpha^\beta \cdot \sqrt{-1_q},$$

and

$$(35) \quad (S q)^2 - (V q)^2 = T q^2 (\cos^2 \alpha^\beta + \sin^2 \alpha^\beta) = T q^2 = (S q)^2 + (T V q)^2.$$

The second of equations (34) is equivalent to the two equations

$$(34') \quad S U q = \cos \alpha^\beta, \quad V U q = \sin \alpha^\beta \cdot \sqrt{-1_q}.$$

The latter of these is equivalent to the two equations

$$(34'') \quad T V U q = \pm \sin \alpha^{\beta*}, \quad U V U q = \pm \sqrt{-1_q}.$$

By the substitution of (34') in (33) and (33a), we obtain

$$(36) \quad S q = T q \cdot S U q, \quad V q = T q \cdot V U q.$$

Finally, by means of (34), (11), and (34'), we get

$$(37) \quad U q^m = (\cos \alpha^\beta + \sin \alpha^\beta \cdot \sqrt{-1_q})^m = \cos m \alpha^\beta + \sin m \alpha^\beta \cdot \sqrt{-1_q},$$

which coincides with the expression of DE MOIVRE'S *Theorem*.

\* Tensors being always positive, if  $\alpha^\beta > 180$ , it would be necessary to use the lower signs.

# NOTES ON THE THEORY OF PROBABILITIES.

By SIMON NEWCOMB, Cambridge, Mass.

[Continued from Page 835, Vol. I.]

15. THE problem which we are about to consider is one of the most fruitful in the theory of probabilities, as out of it grow the theory of errors, the theory of chance distribution, the law of averages, and the estimation of the probability that an observed concurrence of events is the result of a law of nature.

*To find the probability that an event of which the probability on a single trial is  $p$  will happen  $s$  times on  $n$  trials.*

The probability that it will fail on every trial is  $(1 - p)^n$ ,  $1 - p$  being the probability that it will fail on any single trial.

The probability that it will happen on the first trial and fail on the  $n - 1$  following ones is  $p(1 - p)^{n-1}$ . But as the single event is as likely to occur on the 2d, 3d, . . . .  $n$ th trial as on the first, the probability that it will occur just once is  $np(1 - p)^{n-1}$ .

The probability that the event will occur on the first two trials and fail on the  $n - 2$  subsequent ones is  $p^2(1 - p)^{n-2}$ . But the two events can equally occur on the (1, 3), (1, 4) . . . . (1,  $n$ ), or the (2, 3), (2, 4), &c. trials; in fact there will be  $\overset{2}{C}_n$  pairs of trials on which the two events can occur, so that  $\overset{2}{C}_n p^2(1 - p)^{n-2}$  is the probability that it will occur twice.

By a process of reasoning exactly like the last, we find the probability that it will occur  $s$  times to be

$$(1) \quad P_s = \overset{s}{C}_n p^s (1 - p)^{n-s},$$

which is the  $(s + 1)$ st term in the development of the binomial  $[(1 - p) + p]^n$ . The sum of the probabilities of all the possible results of the  $n$  trials is therefore 1, as it ought to be.

As an example to elucidate the above, suppose that a cent is so formed that a head is twice as likely to be thrown as a tail, so that the probability of the former on each throw is  $\frac{2}{3}$ . If the coin is thrown four times, the results of the four throws may be as follows. After each separate result is written the fraction expressing the probability of that result.

No heads,  $tttt \frac{1}{81} = \frac{1}{81}$ ,

1 head,  $httt, thtt, ttht, tthh$ , each  $\frac{2}{81}$ ;  $\times 4 = \frac{8}{81}$ ,

2 heads,  $hhtt, htth, htth, thth, tthh, tthh$ ,  $6 \times \frac{4}{81} = \frac{24}{81}$ ,

3 heads,  $hhht, hthh, hthh, thhh$ ,  $= \frac{24}{81}$ ,

4 heads,  $hhhh = \frac{16}{81}$ .

If we supposed heads as likely to be thrown as tails, we should find these probabilities to be  $\frac{1}{16}, \frac{4}{16}, \frac{6}{16}, \frac{4}{16}, \frac{1}{16}$ , respectively. The result would evidently be the same if we supposed that four coins were thrown from a box together.

16. To resume the general discussion, let us see what value of  $s$  is the most probable. This value we will determine by the condition that its probability must be greater than that of the next smaller number, and also greater than that of the next greater number, or

$${}_nC_n p^s (1-p)^{n-s} > {}_nC_n^{s-1} p^{s-1} (1-p)^{n-s+1};$$

$${}_nC_n p^s (1-p)^{n-s} > {}_nC_n^{s+1} p^{s+1} (1-p)^{n-s-1}.$$

Since  ${}_nC_n = \frac{n-s+1}{s} {}_nC_n^{s-1}; \quad {}_nC_n^{s+1} = \frac{n-s}{s+1} {}_nC_n^s;$

we have from the division of the first inequality by  ${}_nC_n^{s-1} p^{s-1} (1-p)^{n-s+1}$ ,

$$1-p < \frac{n-s+1}{s} p, \quad \text{which gives } s < p(n+1);$$

and from the division of the second inequality by  ${}_nC_n^s p^s (1-p)^{n-s}$  we have

$$1 - p > \frac{n-s}{s+1} p, \quad \text{which gives } s > p(n+1) - 1;$$

$s$  is therefore the greatest whole number in  $p(n+1)$ . If  $s$  and  $n$  are very large numbers, we have very nearly

$$(2) \quad \frac{s}{n} = p.$$

It follows, therefore, that in a great number of trials events are more likely to occur a number of times proportional to their respective probabilities than any other number. Thus if a cent is thrown one hundred times, heads are more likely to be thrown fifty than any other single number of times. But it must not be supposed that they are therefore more likely to be thrown fifty times than not, for it is *almost* as likely to be thrown 49 times or 51 times as fifty times. The chances that it would be thrown *exactly* fifty times would be quite small, because there are so many other numbers that might be thrown.

17. Another deduction from the expression (1) is the following: however small the probability of an event on a single trial, by increasing the number of trials we can render the probability that the event will occur at least once as great as we please. For the probability that it will fail on every one of the  $n$  trials being  $(1-p)^n$ ; however small  $p$  may be, we may make  $n$  so great that  $(1-p)^n$  shall be as small as we please.

18. Suppose now that  $n$  is infinitely great, and  $p$  infinitely small, and that  $np = \alpha$ ,  $\alpha$  being a finite quantity. We may then put  $n = n - 1 = n - 2$ , &c. We shall then have, while  $s$  is finite,

$${}_nC_s p^s = \frac{n^s p^s}{s!} = \frac{\alpha^s}{s!},$$

$$(1-p)^{n-s} = (1-p)^n = (1-p)^{\frac{\alpha}{p}} = e^{-\alpha},$$

$e$  being the Napierian base. Substituting these values in (1) we obtain for the probability that the event will occur  $s$  times

$$(3) \quad P_s = \frac{\alpha^s e^{-\alpha}}{s!}.$$

The probability that the event will fail, is therefore  $e^{-\alpha}$ ; that it will occur once only,  $\alpha e^{-\alpha}$ ; twice only,  $\frac{\alpha^2}{2} e^{-\alpha}$ , &c.

The sum of this series of probabilities continued to infinity is

$$e^{-\alpha} (1 + \alpha + \frac{\alpha^2}{2!} + \frac{\alpha^3}{3!} + \&c.) = e^{-\alpha} \cdot e^{\alpha} = 1, \text{ as it ought to be.}$$

19. We may apply this equation to the determination of the probability that, if the stars were scattered at random over the heavens, any small space selected at random would contain  $s$  stars. Let  $N$  be the whole number of stars,  $h$  the number of units of space in the heavens, then  $\frac{N}{h} dh$  may be taken to represent the infinitely small probability that the infinitely small space  $dh$  contains a star. Moreover, if  $l$  represents the extent of space selected at random which we consider, we may consider the examination of each  $dh$  as a trial, and the number  $n$  of trials will then be  $\frac{l}{dh}$ . The value of  $\alpha$  will then become  $\frac{N}{h} l$ , and by substitution in (3) we have

$$(4) \quad P = \frac{N^s l^s e^{-\frac{Nl}{h}}}{h^s s!}$$

for the probability that the space  $l$  contains  $s$  stars. Suppose, as a numerical example, that  $l$  is a square degree, and  $N = 1500$ , which is about the number of stars of the fifth and higher magnitudes;  $s = 6$ . We then have  $\frac{l}{h} = \frac{1}{41253}$ ; by the substitution of these values in (4) we shall have the probability that any square degree selected at random in the heavens contains six stars. Multiplying this probability by 41253, the number of square degrees in the whole heavens, we obtain the probability that, *if the heavens were divided at random into square degrees, some one of those square degrees would contain six stars.* This probability we find to be

$$\frac{1500^6}{41253^3 \cdot 6!} \cdot e^{-.08636} = .000000128.$$

This, however, is evidently rather smaller than the probability that six stars should be found so near together that a square degree could be fitted on so as to include them.

Mr. MITCHELL committed an error in his solution, the effect of which, if I mistake not, is to make his probability too great. His general method is, however, better applicable to this particular problem than that given above, but as there is a margin of vagueness and uncertainty about the problem in question, so that the answer does not admit of being expressed in exact numbers without an excessively complicated process of reasoning, I have preferred to deduce an approximate solution from the general formulæ, to be used in so many more problems.

20. Let us now consider Prof. FORBES'S objections to the above results of the calculus of probabilities.

He scattered paint from a brush upon a wall, and found double and triple spots and groups innumerable. This is about as decisive as an attempt to disprove the Pythagorean proposition by measuring the squares described on a triangle without knowing whether it had or had not a right angle, and finding that one square was not equal to the sum of the others. As a mathematician would answer this objection by saying that his result only proved that he had either made a mistake in his measurements, or had not measured a right triangle; so Prof. FORBES'S result only proves that either he was mistaken as to the marked character of the groupings, or that the proximity of the components of each group was the effect of their positions being determined *by the action of the same cause*, which is all that the theory of probabilities claims for the Pleiades. The latter supposition is by no means improbable, because a group of spots might be formed by the breaking up of a drop after it had left the brush.



21. Prof. FORBES remarked that an exactly uniform distribution of stars would not be expected as the result of a random distribution. In this he is correct; and as it is an interesting problem, we shall here determine what law a random distribution may be expected to follow. It may appear paradoxical to assert that the results of chance can be expected to follow any law; but such is really the case, and the formula (3) determines the law. As an example, suppose that the heavens are divided into 1500 equal portions, and that 1500 stars are distributed at random, or, to speak with more philosophical accuracy, that the causes which determine the position of each separate star are entirely independent of those which determine the position of any other. Then by reasoning as in § 19 we find  $\alpha = 1$ ; and by formula (3) the probability that a unit of space selected at will contains no star, will be  $\frac{1}{e} = \frac{1}{2.718...}$ \*; one star,  $\frac{1}{e}$ ; two stars,  $\frac{1}{2e}$ ; three stars,  $\frac{1}{2.3e}$ ; &c. If we then select the whole 1500 units we ought to expect the number which would be found to contain the several numbers of stars to be somewhere near 1500 multiplied by the respective probabilities, or

about	$\frac{1500}{e}$	=	552	portions containing no star,
"	$\frac{1500}{e}$	=	552	" " 1 star,
"	$\frac{1500}{2e}$	=	276	" " 2 stars,
"	$\frac{1500}{2.3e}$	=	92	" " 3 stars,
"	$\frac{1500}{4!e}$	=	23	" " 4 stars,
"	$\frac{1500}{5!e}$	=	4(+)	" " 5 stars,
"	$\frac{1500}{6!e}$	=	1	" " 6 stars,

---

\* The acute reader will perceive that the solutions in §§ 19 and 21 are those of a problem slightly (though not materially) different from that actually propounded.

and it would be quite improbable (about 1 chance to 8) that any space would be found to contain more than six stars.

If any one wishes an experimental illustration of the principle let him take a pint of rice, color a hundred grains of it black, mix the black grains thoroughly with the remainder, and stir the mixture till he finds six or eight of the black grains to form a group by themselves. Before this result arrives he will in all probability be willing to admit that should he ever see such a group in such a mixture, he would not believe that it was formed by indiscriminate mixing.

## A SECOND BOOK IN GEOMETRY.



[Continued from Page 104.]

### CHAPTER VI.

#### THE PYTHAGOREAN PROPOSITION.

91. *Theorem.* If a triangle has one side and the adjacent angles equal respectively to a side and the adjacent angles in another triangle, the two triangles are equal. — *Proof.* Let us suppose that, in the triangles  $ABC$  and  $DEF$ , we have the side  $AB$  equal to the side  $DE$ , the angle at  $A$  equal to the angle  $D$ , and that at  $B$  equal to that at  $E$ . Let us imagine the triangle  $DEF$  to be laid upon  $ABC$  in such a manner as to place  $E$  upon  $B$ , and  $D$  upon  $A$ , which can be done because  $AB$  is equal to  $DE$ . Now, as the angle  $A$  is equal to  $D$ , the line  $DF$  will run in the same direction as  $AC$ , and, as it starts from the same point, will coincide with it. Also, since the angle  $B$  is equal to  $E$ , the line  $EF$  will coincide with  $BC$ . Whence, by article 90, the triangles are equal.

92. *Theorem.* The opposite sides of a parallelogram are equal. — *Proof.* Article 90 gives us the only test of geometrical equality. So that in order to prove this theorem we must show that in a parallelogram like  $ABCD$ ,  $AB$  may be made to coincide with  $DC$ , and  $BC$  with  $AD$ . And this would evidently be done if we could show that the triangle  $ABC$  is equal to  $ADC$ . But in these triangles the line  $AC$  is the same, and by article 87 the adjacent angles  $ACB$  and  $CAB$  are equal to the adjacent angles  $CAD$  and  $ACD$ ; whence, by article 91, the two triangles are equal, and  $AD$  is equal to  $BC$ , and  $AB$  equal to  $DC$ .



93. *Axiom.* If one end of a straight line stands still while the other turns round, the end that moves will *begin* to move in a direction at right angles to that of the line itself. Thus if  $AB$  were to begin to turn about the point  $A$ ,  $B$  would *begin* to move either towards  $C$  or towards  $D$ . [If this proposition is not acknowledged as an axiom, the proof is in Select Propositions, Nos. 12–16.]




94. *Theorem.* The angles of a triangle cannot be altered without altering the length of the sides. — *Proof.* If in any triangle, as  $ABC$ , the sides were unchangeable, any alteration of the angles  $A$  and  $B$  would by article 93 make the point  $C$  move in two directions at once, (namely, at right angles to  $AC$ , and at right angles to  $BC$ .) which is impossible, and therefore the angles cannot be altered.

95. *Corollary.* If the three sides of a triangle are respectively equal to the three sides of another triangle, the angles of one must be equal to those of the other, and the equal angles are enclosed in the equal sides.

96. *Theorem.* If the opposite sides of a quadrangle are equal, the quadrangle is a parallelogram. — *Proof.* If in the quadrangle  $ABCD$  (Fig. B.), the sides  $AB$  and  $CD$  are equal, and also the sides  $AD$  and  $BC$  are equal, then, by drawing the diagonal  $AC$ , we have the triangles  $ABC$  and  $ADC$ , composed of equal sides, and, by article 95, the angle  $DAC$  must be equal to the angle  $ACB$ , and the angle  $DCA$  to the angle  $BAC$ ; whence, by article 89, the figure is a parallelogram.

97. *Theorem.* The area of a rectangle is the product of its length by its breadth. — *Proof.*




By drawing lines, at a distance apart equal to the unit of length, parallel to the sides of the rectangle, we shall (arts. 87–89) divide the rectangle into little squares, each of which is a unit of surface. Moreover, these squares are arranged in as many rows as there are units of length in one side of the rectangle, each row containing as many squares as there are units of length in the other side; so that

the whole number of squares is found by multiplying the length of the rectangle by its breadth.

98. *Scholium.* In the above proof it is taken for granted that the sides of the rectangle can be divided into units of length. This can usually be done by taking the units sufficiently short, that is to say, if the lines are not an even number of inches in length, we may take tenths of an inch as the unit; if they are not even tenths, we can divide them into hundredths, or thousandths, or even millionths, of an inch. If after dividing each line into millionths of an inch anything less than the millionth of an inch were left at either end, it would be too small to be taken into consideration. There would be no error, even in reasoning, from neglecting it. For as long as anything is left at the ends of the lines, we can choose smaller units, but as long as the units are of any size at all, our reasoning holds good, and the rectangle is measured by the product of its dimensions.

99. *Theorem.* If the angles of one triangle are equal to those of another triangle, any two sides of one of the triangles have the same ratio to each other as that of the two sides including the same angle in the other triangle. — *Proof.* Let the triangles  $ABC$  and  $DEF$  be equiangular with respect to each other. Place the vertex  $A$  upon the vertex  $D$ , and allow the side  $AB$  to fall upon the side  $DE$ . Since the angles  $A$  and  $D$  are equal, the line  $AC$  will fall upon the line  $DF$ , and since the angles  $C$  and  $F$  are equal, the line  $BC$  will lie parallel to the line  $EF$ .



Let the sides  $AB$  and  $DE$  be divided into units of length,  $Aa$ ,  $aB$ ,  $Bb$ , &c. Through the

points of division draw lines  $ac$ ,  $bd$ , &c., parallel to  $EF$ . Draw also the lines  $ae$ ,  $Bf$ , &c., parallel to  $DF$ . By article 91, the triangles  $Aac$ ,  $aBe$ , &c. are equal. By article 92,  $ae$  is equal to  $cC$ ,  $Bf$  to  $Cd$ , &c. Hence it is easy to see that  $AB$  is composed of the same number of times  $Aa$ , that  $AC$  is of  $Aa$ , and that in like manner  $DE$  is as many times  $Da$ , as  $DF$  is of  $Dc$ . And thus by article 78,  $DE : DF = AB : DC$ , because each of these ratios is equal to  $Ba : ae$ .

100. *Scholium*. If the lines  $AB$  and  $DE$  do not consist of a certain number of times the first unit of length which we have chosen, we may choose a unit small enough to make the remainder small enough to be neglected.

101. *Definitions*. The right angle, right triangle, legs, and hypotenuse, are defined in articles 14 and 17.


102. *Theorem*. The sum of the three angles of a triangle is equivalent to two right angles.

This proposition has been proved in articles 26–31, 34–36, and 57–62.

103. *Corollary*. The sum of the two angles opposite to the legs of a right triangle is equivalent to one right angle.

104. *Corollary*. If an angle opposite a leg in one right triangle is equal to an angle in another right triangle, the two right triangles are equiangular with respect to each other.

105. *Theorem*. If from the vertex of the right angle in a right triangle, a line be drawn at



right angles to the hypotenuse dividing the hypotenuse into two segments, each leg is a mean proportional between the whole hypotenuse and the segment nearest the leg. — *Proof*. Let  $ABC$  be a right triangle with a right angle at  $C$ . Draw  $CF$  at right angles to  $AB$ . The triangle  $BEC$  is right angled at  $E$ , and has an angle at  $B$  equal that at  $B$  in the triangle  $ABC$ . Hence, by article 104, the triangle  $BEC$  has its angles equal to those of  $ABC$ . Hence, by article 99,  $BE : BC = BC : BA$ . In the same way  $AE : AC :: AC : AB$ .

106. *Theorem*. The square on the hypotenuse is equivalent to the sum of the squares on the legs. — *Proof*. Let  $ACB$  (Fig. A.) be a right triangle, with a right angle at  $C$ , and let a square be

drawn on each side. Draw  $CF$  at right angles to  $AB$ . The figure  $BF$  will be a rectangle, because all its angles will be right angles. It will, therefore, be measured by the product of  $BE$  into  $EF$ ; or (since  $EF = BG$  and  $BG = BA$ ), by the product of  $BE \times BA$ . But since  $BC$  is a mean proportional between the lines  $BE$  and  $BA$ , this product is equal to  $BC \times BC$ , which is the measure of the square on  $BC$ . That is, the measure of the rectangle  $BF$  is the same as that of the square on  $BC$ . In the same manner it may be shown that the rectangle  $AF$  is equivalent to the square on  $AC$ . But the sum of these two rectangles is evidently equal to the square on the hypotenuse.

107. In these thirty-one articles I have given you a proof of the Pythagorean proposition in the usual synthetic form. Parts of the proof are not completely filled out, but the omitted steps are so short and easy, that I think you will have no difficulty whatever in supplying them. Do not be satisfied with understanding each of the thirty-one articles, but examine them closely from the 76th to the 106th, and see whether I have introduced anything which is not necessary to the proof of 106. In making this examination, it will be most convenient for you to proceed backward.


These thirty-one articles have been here introduced as lemmas, i. e. preparatory propositions, for demonstrating the Pythagorean proposition. But they are also each one truths worth knowing, and will aid in establishing many theorems that have no connection with a right triangle.

108. Another mode of analyzing this proposition would be suggested by our knowledge of the fact that any triangle is equivalent to half a rectangle of the same base and altitude. I will not lead you through this analysis, but will simply build up for you by synthesis, a

*Second Proof of the Pythagorean Proposition.*

109. *Definitions.* Any side of a triangle or quadrangle may be called its base, and the altitude of the figure is the distance from the base to the most distant vertex of the figure. This distance is measured by a straight line at right angles to the base, and contained between the vertex and the base, prolonged if need be.

110. *Theorem.* Every parallelogram is equivalent to a rectangle of the same base and altitude. — *Proof.* Let  $ABCD$  be a rectangle, and  $ABEF$  a parallelogram having the same base  $AB$ , and the same altitude  $BD$ . It is manifest that if the triangle  $BD F$  by which the parallelogram overlaps the rectangle is equal to the triangle  $A E C$  by which the rectangle overlaps the parallelogram, the two quadrangles are equivalent. But  $A E$  and its adjacent angles are equal to  $B F$  and its adjacent angles, and therefore the triangles are equal (Art. 91), and the quadrangles equivalent.



111. *Theorem.* Every triangle is equivalent to half a rectangle of the same base and altitude. — *Proof.* Let  $A F B$  be a triangle, and  $A B C D$  a rectangle having the same base,  $A B$  and the same altitude  $B D$  (Fig. C). Continue  $C D$  to  $F$  and draw  $A E$  parallel to  $B F$ . The triangle  $A E F$  has its three sides equal to those of  $A B F$ ; the triangles are, therefore, equal to each other (Art. 95); and each is equal to half the parallelogram  $A B E F$ , which is equivalent to the rectangle  $A B C D$ .

112. *Theorem.* The square on the hypotenuse is equivalent to the sum of the squares on the legs. — *Proof.* Having drawn the figure (Fig. A.), as for the former proof, draw the lines  $C' B$ , and  $B' C$ . The triangle  $A B' C$  has the same base  $A B'$ , and the same altitude  $A E$  as the rectangle  $A F$ , and is equivalent to half that rectangle. The triangle  $A B C'$  has the same base  $A C'$  and the same altitude  $A C$  as the square  $C' C$ , and is equivalent to half that square. So that if the triangles  $A B C'$  and  $A B' C$  are equal, the rectangle is equivalent to the square. But these triangles are equal, for if  $A B' C$  were turned about the vertex  $A$  as on a pivot until the point  $C$  covered  $C'$ , then  $B'$  would cover  $B$ , and the triangles would coincide. For  $A C$  would rotate through a right angle, and  $A B'$  through a right angle; and  $A C = A C'$ , and  $A B = A B'$ .

113. This proof of the Pythagorean proposition is more strictly geometrical than the preceding, as it does not involve the idea of multiplying lines to measure areas. But you must remember that each is equally conclusive. I have here also omitted some of the shorter steps. You should not only be able to fill out these steps when the omission is pointed out to you, but also to discover the omission for yourselves. Take the proofs which I have written down and examine them step by step, asking at each step, is that strictly self-evident? Can it be questioned? Can it be divided into two steps? Is there need of proof? If so, has the proof been given in a previous article? It is only by such an earnest study of the book and of the subject that you can make the process of mathematical reasoning become a sure and pleasant road for you to the discovery of truth.

[To be continued.]

## Editorial Items.

THE following students have sent us solutions of the Prize Problems in the October Number, Vol. II., of the Monthly.

- MISS AMANDA BENNETT, Red Wing, Minnesota. Prob. I.  
MISS HARRIET L. ENSIGN, Catskill Academy, N. Y. Probs. I., II., III., and IV.  
MISS HARRIET S. HAZELTINE, Worcester, Mass. Probs. I., II., and IV.  
OAKLEY H. SMITH, New Hampton Institution, Fairfax, Vt. Probs. I. and II.  
D. Y. BINGHAM, Ellicottville, N. Y. Prob. III.  
SAMUEL S. EASTWOOD, High School, Saxonville, Mass. Probs. I. and II.  
C. P. MARSTON, Public High School, Hartford, Ct. Probs. I. and II.  
WILLIAM W. JOHNSON, Sophomore Class, Yale College, Ct. Probs. III. and IV., and first part of V.  
BENJAMIN F. WEBBER, Wesleyan Seminary, Kent's Hill, Maine. Probs. I. and II.  
ISAAC H. TURRELL, Drewsburg, Ind. Probs. I., II., III., and IV.  
GEORGE D. HALE, Select School, Adams Centre, N. Y. Probs. I. and II.  
F. E. HASTINGS, Literary Institution, Suffield, Ct. Prob. I.  
G. W. BROWN and ISAAC FULLER, Collegiate Institution, Bowden, Ga. Each Probs. I., II.  
WILLIAM C. CLEVELAND, Lawrence Scientific School, Cambridge, Mass. All but V.  
WILLIAM C. HENCK, High School, Dedham, Mass. Probs. I. and II.  
CHARLES B. BOUTELLE, Waterville Academy, Maine. Prob. III.  
M. H. DOOLITTLE, Sophomore Class, Antioch College, Yellow Springs, Ohio. Probs. III., IV., and first part of V.  
W. D. MERSHON, Sophomore Class, Princeton College, N. J. Prob. IV.  
R. E. LEONARD, Ward School, No. 32, New York City. Probs. I. and II.  
J. B. FOSSETT, New London Institute, Ct. Probs. I., II., and III.  
SAMUEL I. BALDWIN, Chester Institute, Chester, N. J. Probs. I. and II.  
S. W. BURNHAM, New York City. Probs. I. and II.  
M. L. STREATOR, Mayslick Academy, Ky. Probs. I., II., and III.  
G. H. TOWER, Classical High School, Petersham, Mass. Probs. I. and II.  
PERRIN B. PAGE, Literary Institute, Nunda, N. Y. Probs. I. and II.  
J. D. VAN BUREN, Rensselaer Polytechnic Institute, Troy, N. Y. Probs. III., IV., and V.  
O. B. WHEELER, Sophomore Class, University of Michigan, Ann Arbor. Probs. III., IV., and V.  
JOSEPH COCKE and AYLETT B. COLEMAN, Locust Grove Academy, Albemarle Co., Va. Each Probs. I. and II.  
M. K. BOSWORTH, Sophomore Class, Marietta College, Ohio. Probs. III., IV., and V.  
W. A. AKEN, New Wilmington, Pa. Probs. I. and II.

The following solutions, unfortunately, did not reach us in time.

WILSON BERRYMAN and OTHO E. MICHAELIS, Sophomore Class, N. Y. Free Academy, each answered all the questions.

H. TIEMAN, Baltimore, Md., answered all the questions but the second part of Prob. V.

PHILO HOLCOMB, Hughes High School, Cincinnati, answered Prob. IV.



# The New York Observer,

## THE LARGEST NEWSPAPER IN THE WORLD.

### NATIONAL, CONSERVATIVE, AND RELIGIOUS.

#### BELONGING TO NO PARTY IN POLITICS, AND TO NO SECT IN RELIGION.

Edited by a corps of Clergymen and Laymen of large experience, having the most eminent writers of the day among its regular contributors, and a Foreign Correspondence unrivalled. It is the most

#### COMPLETE FAMILY NEWSPAPER

that can be made; published on a large double sheet, so that it may be separated, making

#### TWO DISTINCT PAPERS,

each perfect in itself. No other newspaper is made upon this plan.

The Secular Sheet contains a full report of all the news of the day; a vast amount of miscellaneous reading, poetry and prose; an AGRICULTURAL page, conducted by a practical and educated agriculturist; a COMMERCIAL page, edited by a gentleman distinguished for his acquaintance with the financial world, giving the latest reports of the money, produce, and cattle markets, bank stocks, &c.; a MISCELLANEOUS department, embracing scientific, literary, and artistic matter, with tales, anecdotes, biography, travels, questions and answers, for the instruction and amusement of the family and social circle.

The Religious Paper is filled with the choicest original and selected matter in every department of Christian literature, making a delightful Sabbath companion, and furnishing a volume of interesting and instructive reading every week. The best and most accomplished Clergymen, Presidents and Professors in our colleges and seminaries constantly contribute to its pages. One of its chief features of attraction is a summary of intelligence from

#### ALL RELIGIOUS DENOMINATIONS;

a feature peculiar to the OBSERVER, and highly valued by Christians who wish to know what is doing in other communions than their own.

The grand object of the NEW YORK OBSERVER is to promote "peace on earth and good-will among men." For this end it seeks to advance all those principles which make the Union of the States more firm and permanent; it cultivates harmony and good feeling among all denominations of

Christians; and is a fearless defender of the rights of all men, under the Constitution of the United States and Word of God.

In its editorial discussion, its foreign and domestic correspondence, the vigor and beauty of its original contributions, and the attractions of its several departments in science, literature, art, agriculture, and commerce, the NEW YORK OBSERVER is determined not to be surpassed by any newspaper.

Resisting radicalism in Church and State, promoting revivals of pure religion and every wholesome moral reform, on Scriptural and rational principles, discarding and opposing all schisms, humbugs, fanaticism, and every scheme of infidelity, socialism, and vice, the NEW YORK OBSERVER designs to be a safeguard of virtue, law, and order, a champion of truth and righteousness in the earth.

It is the cheapest newspaper of its class that is published. Both the secular and religious papers are sent for Two Dollars and Fifty Cents, in advance. Two families using in taking it, as many do, will each get a complete newspaper for \$1.25.

#### Proposals for Soliciting Subscribers.

To any one who will obtain new subscribers for us, we will pay the following liberal commissions: For five new subscribers, paying in advance, 50 cents each; for more than five and less than ten, 75 cents each; for ten or more \$1 each. We will send a copy of our Bible Atlas, with colored maps, on paper of large size and best quality, to each new subscriber, on the receipt of his name and payment for one year.

If you cannot give personal attention to this work, will you show this advertisement to some clergymen or layman who will take an interest in it, to whom we will give the commissions mentioned above.

We will send specimen numbers without charge.

Your early attention is solicited to this subject, and we shall be happy to hear from you immediately, as we desire to offer the paper at once to every family in the United States.

SIDNEY E. MORSE, JR., & CO.,

Editors and Proprietors, No. 37 Park Row, New York.

#### JUST PUBLISHED.

## Hitchcock's Anatomy and Physiology.

### FOR COLLEGES, ACADEMIES, AND SCHOOLS.

By PRES. HITCHCOCK, D.D., LL.D., of Amherst College, Author of a "Treatise on Geology," etc., etc., and EDWARD HITCHCOCK, JR., M.D., of Williston Seminary, Easthampton, Mass. Illustrated by nearly 400 engravings. 441 pages. Retail price, \$1.00.

THE Publishers have the pleasure of announcing that this work, which has been in preparation for two years past, is now ready; and offer it in entire confidence that it will receive a cordial welcome. Pres. HITCHCOCK is an eminent and successful author of scientific works, a naturalist widely known for his attainments, and has been a distinguished educator for more than forty years. His son has for ten years past been engaged in the work of teaching. Their success as practical instructors, and the conscientious fidelity with which the present volume has been prepared, naturally afford abundant assurance of a superior work. The authors have availed themselves of the labors of the most eminent Physiologists, American and foreign; and they give a condensed but clear exhibition of the leading principles and facts which are detailed in such works as Carpenter's Human Physiology, his Comparative Physiology, Hanall's Microscopic Anatomy, Griffith and Henfrey's Micrographic Dictionary, Peaslee's Histology, Todd and Bowman's Physiological Anatomy, Wilson's Anatomy, Draper's, Dungenion's, and Dalton's Physiologies, and the works of Cuvier, Owen, and Weyner on Comparative Anatomy. The illustrations of the volume are unusually full, were executed by superior artists, French and German, as well as English and American; and their first cost was not less than \$3,000. It is believed that no school-book yet issued has surpassed this in the value of its illustrations, and the important aid that they render in fully understanding the text.

Of several important new features of this book, one is the introduction of **Microscopic Anatomy**, a branch of the subject which has become indispensable in this study, even to the beginner. **Comparative Anatomy and Physiology** are also much more fully treated of here than in any other school-book on this subject now extant.

The introduction of this department of Anatomy and Physiology it is

hoped will not only give some knowledge of the organs and their use in the lower animals, but will at the same time give a much better means of ascertaining the structure and functions of the human mechanism. Another new feature of the work is the religious bearing of the subject, in a chapter written by the senior author. The few technical terms which it has been found indispensable necessary to introduce are so fully explained that the simplest intellect cannot fail at once, and readily, to understand them.

Considerable attention has also been given to the important subjects of **Hygiene and Gymnastics**; the latter is illustrated with diagrams, designed to facilitate the practical use of these means of healthful and pleasant exercise in schools and families. **Ventilation, bathing, and other personal habits**, bearing upon the preservation of health, and many other subjects of practical usefulness, as well as scientific interest, are also fully discussed and illustrated. The moral tone of the work in every respect is unexceptionable; the utmost care having been taken to treat the subjects not merely with delicacy, but with appropriate reference to the wisdom and goodness of the Creator.

The *Springfield Republican* says:—"At length we have the prospect of a class-book of Anatomy and Physiology thoroughly adapted to the use of schools. We have had numerous books on these comparatively new school studies, but they do not live. Their sufficient condemnation is that they die out. Pres. HITCHCOCK, of Amherst College, and his son, Dr. HITCHCOCK, of Williston Seminary, have prepared a book, and from what we know of the authors, we cannot doubt that it will sweep the market. The work is handsomely illustrated, and especial attention given to Gymnastics and Hygiene. The men not only understand their subject, but are teachers, and know how to adapt their instructions to the purposes of schools."

Single Copies to TEACHERS for examination, with a view to introduction into classes, will be furnished at our store, or sent, prepaid, on receipt of 50 cents.

Published by IVISON & PHINNEY,

48 and 50 Walker Street, New York.

SOLD BY BOOKSELLERS GENERALLY.



# WEST POINT COURSE OF MATHEMATICS.

STUDENTS entering the *Military Academy of the United States* at West Point are examined in THE UNIVERSITY ARITHMETIC, by Charles Davies, LL.D., Professor of Mathematics in Columbia College, New York, and Author of a Complete Course of Mathematics, for Schools, Academies, and Colleges. The following are the TEXT-BOOKS used in the Institution, by Professors Church and Bartlett.

Davies's Bourdon's Algebra.  
 Davies's Legendre's Geometry and Trigonometry.  
 Davies's Elements of Surveying.  
 Davies's Descriptive Geometry.  
 Davies's Shade, Shadows, and Linear Perspective.  
 Church's Analytical Geometry.  
 Church's Elements of Differential and Integral Calculus.  
 Bartlett's Analytical Mechanics.  
 Bartlett's Acoustics and Optics.  
 Bartlett's Spherical Astronomy.

P.S.—The above Text-Books (with one or two exceptions) are the STANDARD in the FREE ACADEMY of New York City, the ROCHESTER UNIVERSITY, UNIVERSITY OF VIRGINIA, and many other Institutions of Learning.

Published by A. S. BARNES & BURR,

51 and 53 John Street, New York.

## LIST OF NEW PUBLICATIONS

ISSUED BY

BROWN, TAGGARD, & CHASE,

25 and 29 Cornhill, Boston.

### Miscellaneous.

STRICKLAND'S QUEENS OF ENGLAND. The illustrated edition. Seven volumes, 8vo. Price, \$1.75 per volume.

MRS. CORNELIUS'S COOK-BOOK. One volume. 12mo. Price, 75 cents.

AN ARCTIC BOAT-JOURNEY. By Dr. I. I. HAYES, Surgeon of the Kane Expedition. (To be ready about January 1st.) One volume. 12mo. \$1.00.

POETS AND POETRY OF VERMONT. One volume. 12mo. Price, \$1.25.

ARCTIC ADVENTURES. By EPES SARGENT. A new edition. Price, \$1.25.

LIFE WITHOUT AND LIFE WITHIN. By the late MARGARET FULLER; with a new steel portrait of the authoress. (Ready about January 1st.) One volume. 12mo. Price, \$1.25.

At the same time, new and uniform editions of her previous works, comprising,

AT HOME AND ABROAD.

WOMAN OF THE NINETEENTH CENTURY.

ART, LITERATURE, AND THE DRAMA.

Together with

THE MEMOIR OF MARGARET FULLER. In two volumes.

### Educational.

EATON'S ARITHMETIC. Price, 75 cents.

UNIVERSAL SPEAKER. By N. A. CALKINS and W. T. ADAMS. Price, \$1.00.

WELD AND QUACKENBOS'S NEW PROGRESSIVE ENGLISH GRAMMAR. Price, 60 cents.

PRIMARY SCHOOL TABLETS. 20 in number. Each card consists of two Tablets. Prepared by JOHN D. PHILBRICK, Superintendent of the Public Schools of Boston. Price, 50 cents per card.

### Juvenile.

OUR SUMMER-HOUSE. By the Author of "Daisy," "Violet," &c. Beautifully illustrated. Price, 62 cents.

NELLIE AND HER FRIENDS. Beautifully illustrated. Price, 62 cents.

PICTURES FROM THE HISTORY OF THE SWISS. Beautifully illustrated, and very attractive. Price, 75 cents.

LIFE AND ADVENTURES OF WHITENOSE WOODCHUCK. Designed for younger children, and splendidly illustrated. Price, 38 cents.

These four books are entirely new and original. The first two are the first volumes of new series, entitled, "The Summer-house Series," and "The Martin and Nellie Stories," so written, however, that each book is distinct in itself.

### Our Standard Juveniles are

ROLLO'S TOUR IN EUROPE. In ten volumes, put up in a nice box. Price per set, \$5.00.

MY UNCLE TOBY'S LIBRARY. In twelve volumes. In boxes. Price per set, \$3.00.

### New Music-Book.

THE CHORAL HARMONY. By B. F. BAKER and W. O. PERKINS. Price, \$1.00.

The only new music-book issued this season. The demand is so great as to convince the publishers that it will be very extensively and permanently used.

# WELLS'S NATURAL PHILOSOPHY.

For Schools, Colleges, and Families.

450 pages; 374 illustrations. Twentieth Edition. Retail Price, \$1.00.

FOR USE WITH OR WITHOUT APPARATUS.

From among the hundreds of Testimonials from eminent and well-known educators, the following are offered as specimens. For several pages more of similar character, and for a list of HUNDREDS OF THE BEST ACADEMIES, &c. in this country, where the work is used with high success and approval, see IVISON, PHINNEY, & Co.'s QUARTERLY EDUCATIONAL NEWS, and their NEW DESCRIPTIVE CATALOGUE (of 172 pages) both sent free on request.

Prof. WILLIAM HOPKINS, A.M., Genesee College, New York, writes, March 17, 1858:—"I find 'Wells's Philosophy' a Work of superior merit, either for the student or general reader. The arrangement and style I consider excellent."

THOMAS SHERWIN, A.M., Principal of the Boston City Normal School, writes:—"In examining 'Wells's Philosophy' I have found it more nearly correct, and more up to the times, than text-books in general use."

E. CONANT, A.M., Principal of Normal Institute, Royalton, Vt., writes:—"I have made use of 'Wells's Philosophy' for several terms, and find it the most satisfactory text-book on the subject that I know of."

Rev. J. R. BOYD, A.M., Principal Maplewood Female Institute, Pittsfield, Mass.:—"We have adopted 'Wells's Philosophy,' and are greatly pleased with it."

Prof. EBENEZER EMMONS, Professor of Natural Sciences, Williams College, State Zoologist, N. C.:—"Wells's Series of text-books have my highest recommendation."

Rev. D. C. VAN NORMAN, D.D., Principal of Young Ladies Institute, New York:—"To the practical teacher, a cursory examination of 'Wells's Philosophy' will, I think, suffice to show its superiority over any school-book of the kind."

H. M. PIERCE, A.M., Principal Rutgers Female Institute, New York:—"We have used 'Wells's Philosophy' and 'Chemistry' for some time past, with the highest satisfaction."

Prof. CLEVELAND, Philadelphia:—"I am very much pleased with 'Wells's Philosophy,' and shall introduce it."

ALFRED BROOKS, M.D., Lecturer on Physics and Chemistry, Southwick, Mass.:—"I consider 'Wells's Philosophy' far superior to any other work of like grade now before the public. The exceptions of a New York critic (Harper's Magazine) to some portions of it appear to me to be frivolous, and to indicate personal feeling rather than fair criticism. We do not want, in an elementary work, statements of every exception to general rules; but a clear outline of the great leading facts and principles; and this we find most perfectly in Wells's Scientific Series."

AVERY S. WALKER, Esq., Superintendent of Public Schools, Lodi, N. J.:—"After careful examination, I deem 'Wells's Philosophy' a work of much merit, and well adapted to teaching the science."

C. F. WINSLOW, M.D., an eminent Physicist:—"Wells's Series of Elementary Scientific Text-books pleases me more than any other works of like character which I have ever examined; and I believe I do good service to the cause of science and education in most unreservedly recommending them."

N. T. CLARK, A.M., Principal of Canadaigua Academy, N. Y.:—"We are using 'Wells's Philosophy' with great satisfaction. It is very much superior to Parker's, and better suited to our wants than Olmsted's."

Pres. A. W. CUMMINGS, D.D., President of Holston Conference Female College, N. C., writes, June, 1859:—"We have used 'Wells's

Philosophy,' and the least I can say of it is, that we regard it as a great advance upon all former efforts to present the whole science of Natural Philosophy in an easy yet strictly scientific and perspicuous manner. It is destined to be a highly popular text-book."

ANDREW J. RICKOFF, Esq., Superintendent of Schools in Cincinnati, and an eminent educationist, writes, May, 1859:—"My examination of Wells's works has given me unusual pleasure, because they seem to approximate more nearly to the true ideal of a perfect elementary school-book than any others with which I am acquainted, whether upon these or any other school studies. I have no hesitation, therefore, in saying that *I believe they have no equals in their respective departments.*"

Prof. J. H. WILLARD, Principal of the Troy Female Seminary, has also recently introduced "Wells's Philosophy" and "Chemistry" into that important institution.

Prof. E. W. EVANS, Professor of Natural Philosophy, Marietta College, Ohio, writes:—"The 'Philosophy' is the most comprehensive, the best illustrated, manual for schools and academies that I have ever seen on the subject."

Prof. J. SANDFORD BARNES, United States Naval Academy, Annapolis, Md., writes:—"It is without hesitation, as it is with great pleasure, that I say it is, in my opinion, better adapted to popular instruction than any work of the kind ever published in this country."

HORACE WEBSTER, LL.D., Principal New York Free Academy, writes:—"Wells's Philosophy' is the best work on this subject, of its grade, that I ever saw."


Rev. CHARLES COLLINS, D.D., President of Dickinson College, writes:—"Wells's Philosophy' I consider the best elementary treatise on that subject now before the public."

Prof. DANIEL VAUGHAN, an eminent author and lecturer on Physics, in Cincinnati, writes:—"I have been much delighted with Wells's valuable works on 'Natural Philosophy' and 'Chemistry.' I am acquainted with no scientific books better adapted for schools and seminaries. The 'Philosophy' must be regarded as a great boon in the cause of education, as the works in this department generally studied are very defective."

Hon. H. C. HICKOK, Superintendent of Public Instruction of the State of Pennsylvania, whose critical ability and care are well known, writes:—"A cursory examination of Wells's admirably illustrated and arranged works impress me with the belief that they are well adapted to their purpose, and would be invaluable in the school and family circle."

His Excellency JAMES POLLOCK, Ex-Governor of Pennsylvania:—"I have examined carefully and in detail this series of books by Mr. Wells, and am more than satisfied with their value and importance as auxiliaries in the great cause of true and practical education. The general principles of the sciences of which they treat are stated clearly and concisely, and the latest results of scientific discovery and research are fully and correctly embraced in these volumes. They are admirably adapted to meet the wants of students in our schools, academies, and colleges, and they should have a place in the library of every family in the country."

Another page of similar notices may appear in the next number; also of commendatory reviews from the principal STATE EDUCATIONAL JOURNALS, and other LITERARY AND SCIENTIFIC PAPERS.

 To enable Teachers to verify, by personal examination, the truth of these commendations, single copies will be sent by mail, prepaid, on receipt of one half the published price.

ESPECIALLY LIBERAL TERMS FOR FIRST INTRODUCTIONS.

IVISON, PHINNEY, & CO., Publishers, New York.

Sold by Booksellers generally.

A LIBERAL OFFER CONTINUED.

## DAVIES & PECK'S MATHEMATICAL DICTIONARY.

In consequence of the continued and repeated calls for the "DICTIONARY OF MATHEMATICS," by DAVIES AND PECK, the publishers have decided to extend their offer to all Teachers of Mathematics, (whether Prof. Davies's course, or any other,) to send them a copy of this sterling work, *post-paid*, for ONE DOLLAR AND THIRTY-NINE CENTS, until the 1st of February, 1860; after which time the price will be as before, viz. \$2.50.

The very high commendations of this work from the most distinguished sources warrant the publishers in urging the attention of all teachers of the science to this opportunity of securing a superb work at a very low price.

A. S. BARNES AND BURR,

PUBLISHERS OF

### DAVIES'S COMPLETE COURSE OF MATHEMATICS.

To this popular series of Text-books have recently been added two entirely new works on Algebra, entitled

**NEW ELEMENTARY ALGEBRA;** embracing the First Principles of the Science. By CHARLES DAVIES. 299 pp. 12mo. Cloth. Price, 75 cents.

BLOOMINGTON, ILL., Sept. 5, 1859.

Messrs. A. S. BARNES AND BURR:—

Gentlemen:—I have examined Davies's New Elementary Algebra, and am free to say that I deem it superior in every respect to any work on Elementary Algebra that has come under my observation.

Very respectfully yours,

JOSEPH FICKLIN,

Prof. Mathematics in Bloomington Female College.

**UNIVERSITY ALGEBRA;** embracing a Logical Development of the Science, with numerous Graded Examples. By CHARLES DAVIES. 320 pp. 12mo. Sheep. Price, \$1.00.

SYDNEY, O., Sept. 30, 1859.

SIRS:—I received a copy of the University Algebra, and from a cursory examination of it, I think it is destined to supersede all other Algebras above the elementary grade in our Union Schools and Seminaries, as it comprises, in a convenient, concise, and compact form, all that is usually required in that branch of mathematics.

Yours, &c.,

W. H. SCHUYLER,

Principal of Union School.

#### Just issued likewise:

**ELEMENTS OF MECHANICS.** For the Use of Colleges, Academies, and High Schools. By WM. G. PECK, M. A., Professor of Mathematics, Columbia College. 338 pp. 12mo. Cloth. Price, \$1.25.

Prof. B. F. STEM, Principal of Easton (Pa.) High School, writes:—

"This book supplies a want long felt. It is a work of deep research, simple in its arrangement, complete in its examples and rules, and is an admirable volume to assist the intelligent teacher. There can be no doubt but that it will have a large sale, as it deserves to have."

Copies of these new works will be furnished teachers for examination, post-paid, on receipt of one half their respective prices.

A. S. BARNES & BURR, Publishers.

**NATURAL PHILOSOPHY FOR COLLEGES.** By W. H. C. BARTLETT, LL. D., Professor of Natural Philosophy in the Military Academy of the United States.

BARTLETT'S SYNTHETIC MECHANICS . . .	\$3.00
BARTLETT'S ACOUSTICS AND OPTICS . . .	2.00
BARTLETT'S SPHERICAL ASTRONOMY . . .	3.50
BARTLETT'S ANALYTICAL MECHANICS . . .	4.00

From JOHN A. NICHOLS, Professor of Natural Philosophy in the Free Academy, New York:—

"Professor Bartlett's Course of Natural Philosophy, entitled Mechanics, and Acoustics, and Optics, have been text-books in the department of Natural Philosophy in this institution about two years. They have been found, by a daily test in the class-room, to possess that essential quality of works of merit, namely, the more familiarly they are known, the better they are liked. They develop the principles of Natural Philosophy by means of the Mathematical Analysis: and, by the aid of the same unflinching agent, investigate the laws of machinery, the effects of friction, and other resisting forces, and the motion of the planets, basing every process upon the fundamental results of observation and experiment.

"The methods of Professor Bartlett's works reward without discouraging effort; and, when faithfully studied, these works give a discipline of mind of the highest importance to any man, whether he seeks for reputation and support in the study, the workshop, the mart, or the forum."

**CHEMISTRY FOR COLLEGES.** By WM. GREGORY, M. D., F. R. S. E., Professor of Chemistry in the University of Edinburgh. Edited by J. M. SANDERS, M. D., LL. D.

**HAND-BOOK OF INORGANIC CHEMISTRY.** For the Use of Students. To which is added the Physics of Chemistry. 426 pp. 8vo. Cloth. Price, \$1.50.

**HAND-BOOK OF ORGANIC CHEMISTRY.** For the Use of Students. 480 pp. 8vo. Cloth. Price, \$1.60.

"Among the excellences of these works are a clear statement of facts and principles, a concise mode of explanation, and a scientific arrangement. Together they form one of the best manuals extant in chemical science. They contain all the 'modern improvements' that have attained the dignity of scientific truth. We give them our hearty commendation."—*New York Teacher*.

Among the recent additions to the National Series of Standard School-Books, published by A. S. BARNES AND BURR, are several very valuable ones to the popular

#### SCHOOL-TEACHER'S LIBRARY.

**THE NORMAL;** or, Methods of Teaching the Common Branches, Orthoepey, Orthography, Grammar, Geography, Arithmetic, and Elocution. By ALFRED HOLBROOK, Principal of Normal School, Lebanon, Ohio. 456 pp. 12mo. Cloth. Price, \$1.00.

"This book is designed to bring to the teacher's view a model working school. It is not all theory, but theory and practice combined. It is a text-book on the science of teaching, a sort of perpetual Institute to every teacher who is so fortunate as to possess a copy. Those of the largest experience will here find things new, while the merest tyro in teaching, with 'The Normal' for an instructor, need never be at fault in his school."—*School Visitor*.

**THE HIGHER CHRISTIAN EDUCATION.** By BENJAMIN W. DWIGHT. 347 pp. 12mo. Cloth. Price, \$1.00.

From CHARLES NORTHEND, Author of "Teacher and Parent."

"It is a work of rare merit. The book will do good. I shall earnestly commend it."

**INSTITUTE LECTURES ON MENTAL AND MORAL CULTURE.** By SAMUEL P. BATES, A. M., Supt. of Public Instruction, Crawford County, Pa. 319 pp. 12mo. Cloth. \$1.00.

"We find these Lectures well written, and replete with sound views and full information."—*Philadelphia Press*.

The TEACHER'S LIBRARY now includes the following volumes, all of incalculable value to the growing teacher:—

PAGE'S THEORY AND PRACTICE OF TEACHING . . .	\$1.00
NORTHEND'S TEACHER AND PARENT . . .	1.00
ROOT ON SCHOOL AMUSEMENTS, GYMNASICS, &c. . .	1.00
MANSFIELD ON AMERICAN EDUCATION . . .	1.00
MAYHEW ON UNIVERSAL EDUCATION . . .	1.00
DE TOCQUEVILLE ON AMERICAN INSTITUTIONS . . .	1.00
DAVIES'S LOGIC OF MATHEMATICS . . .	1.00
HOLBROOK'S NORMAL METHODS OF TEACHING . . .	1.00
DWIGHT'S HIGHER CHRISTIAN EDUCATION . . .	1.00
BATES'S INSTITUTE LECTURES . . .	1.00
BARNARD ON SCHOOL-HOUSE ARCHITECTURE . . .	2.00
BARNARD ON NORMAL SCHOOLS . . .	2.00

Of these volumes, and about two hundred other valuable educational works, a complete and extended description may be found in the publisher's Illustrated Descriptive Catalogue, which will be forwarded to any address on application.

A. S. BARNES AND BURR, 51 and 53 John Street, New York.

# Popular and Valuable Educational Publications

OF

IVISON, PHINNEY, AND COMPANY,  
NEW YORK.

## FASQUELLE'S FRENCH TEXT-BOOKS.

By L. FASQUELLE, LL. D.

I. FRENCH COURSE, 70th edition, . . . . .	\$ 1.25
II. KEY TO FRENCH COURSE, . . . . .	.63
III. JUVENILE FRENCH COURSE, . . . . .	.56
IV. COLLOQUIAL FRENCH READER, . . . . .	.75
V. TELEMACHUS, . . . . .	.75
VI. DUMAS'S NAPOLEON, . . . . .	.75
VII. CHEFS D'ŒUVRE DE RACINE, . . . . .	.75
VIII. MANUAL OF FRENCH CONVERSATION, . . . . .	.75
IX. HOWARD'S AIDS, . . . . .	1.00
X. TALBOT'S FRENCH PRONUNCIATION, . . . . .	.75

The plan of this popular Series embraces a combination of two rival systems; the *Oral*, adopted by Ollendoff, Robertson, Manesca, and others, with the old Classical or *Grammatical System*.

This Series is now in use in a large majority of the best Colleges and Academies in New England, having been adopted in hundreds of

those Institutions during the past year. Of the one hundred and seventy-four incorporated Academies in the State of New York, in which the French language is taught, *Fasquelle's Series* is used in over one hundred, whilst the highest number of Institutions using any other series, or single volumes, is fifty-three. It is also the leading French Series in New Jersey, the Northwestern States, and in many others; and some of the volumes have been republished in England, where they have enjoyed a very extensive sale for several years. These facts form the safest expression of the practical value and adaptation of this Series that could be given.

In our *Descriptive Catalogue*, and previous numbers of the *Educational News*, we present strong recommendations from hundreds of practised and successful Teachers of the French Language, from all parts of the Union.

## WOODBURY'S GERMAN SERIES.

By W. H. WOODBURY, A. M.

I. NEW METHOD WITH GERMAN, . . . . .	\$ 1.50
II. KEY TO THE NEW METHOD, . . . . .	.50
III. SHORTER COURSE, . . . . .	.75
IV. KEY TO SHORTER COURSE, . . . . .	.50
V. ELEMENTARY GERMAN READER, . . . . .	.75
VI. ECLECTIC GERMAN READER, . . . . .	1.00
VII. GERMAN ENGLISH READER, . . . . .	.25
VIII. NEW METHOD FOR GERMANS, . . . . .	.75

Prof. Hoshour, N. W. C. College.  
Prof. Perry, Williams College.  
Prof. Schmidt, Columbia College.  
Prof. Baermann, Troy Female Seminary.

Prof. Picard, Illinois College.  
Prof. Torricelli, Dartmouth College.  
Prof. Miller, Hartwick Theological Sem.  
Prof. Piessner, Union College.

IX. GLAUBENSKLEE'S GRAMMAR, . . . . .	\$ 0.75
X. GLAUBENSKLEE'S READER, . . . . .	.50

This system is founded on similar principles with *Fasquelle's* French Series, and is highly popular. Reference to our *Catalogue*, and previous numbers of the *Educational News*, presents cordial testimonials from a large number of eminent educators and scholars, among whom are

Pres. Allen, Girard College.  
Prof. Fasquelle, University of Michigan.  
Prof. Jarman, Union University, Tenn.  
Rev. Dr. Nast, Cincinnati.

Single Copies of the above, such as are used in Classes, sent at half price to Teachers for examination, with a view to Introduction. Liberal terms for first supplies.

## THE AMERICAN DEBATER;

BEING A

Plain Exposition of the Principles and Practice of Public Debate.

By JAS. N. McELIGOTT, LL. D.,

PRESIDENT OF THE NEW YORK STATE TEACHERS' ASSOCIATION.

324 pages. Price, \$ 1.00.

This book presents a mass of information respecting debates and debaters not to be found in any other single volume. It carefully, yet briefly, shows the qualifications essential to a good debater, and the best modes of acquiring them; furnishes a full course of instruction in the Rules of Parliamentary Practice, grounded on the best authorities, and illustrates the practical working of an orderly debate by giving debates in full on interesting questions. It furnishes Debates in Outline, to be filled out according to the taste and ability of the parties that use them; presents a series of questions, with references under each, to authorities or sources of information; supplies nearly six hundred interesting questions for discussion, and gives forms of a Constitution and By-Laws for Debating Societies.

The following journals have commended the Debater:—

New York Tribune, Boston Traveller, Chicago Daily Journal, New York Chronicle, Albany Express, Richmond Post, Springfield Republican, New York Evening Post, Knickerbocker Magazine, Chicago Press, Boston Post, New York Albion, Chicago Christian Times, Congregational Journal, Christian Herald, R. I. Schoolmaster, New York Observer, Boston Atlas, New York Evangelist, Commercial Advertiser, Newark Whig, Life Illustrated, New York Daily News, New York Day-Book, &c., &c.

Sent, post-paid, on receipt of the price, \$ 1.00.

IVISON, PHINNEY, & CO., Publishers, New York.



HARPER & BROTHERS will send either of the following Works by Mail, postage paid (for any distance in the United States under 3,000 miles), on receipt of the Money.

## ELEMENTS OF NATURAL PHILOSOPHY, DESIGNED FOR ACADEMIES AND HIGH SCHOOLS. BY ELIAS LOOMIS, LL.D.

12mo. 350 pages. Sheep extra. \$ 1.00.

This volume is the result of many years of laborious study, and presents in an elementary form the fruits of much research. It is designed particularly for the use of the higher classes in Academies and High Schools, and its primary object is to cultivate the habit of philosophical reflection and observation.

The following are a few of the Notices of LOOMIS'S NATURAL PHILOSOPHY which have been received by the Publishers:—

"Professor Loomis's Elements of Natural Philosophy is one of the best and most complete works of the kind that has been published, and possesses a clearness and conciseness seldom equalled. I have seldom seen so much useful information in one volume."—EDWARD T. FRISTOE, *Professor of Mathematics, Columbian College, D. C.*

"This volume has the merit of being a digest, and not a compilation. For a large class of students it is the only suitable work on this subject that I have seen, bringing into use, as it does, all the mathematics usually studied in our academies, and no more."—E. W. EVANS, *Professor of Natural Philosophy in Marietta College, Ohio.*

"Professor Loomis's Natural Philosophy is clear, accurate, and methodical."—DANIEL KIRKWOOD, *Professor of Natural Philosophy in Indiana State University.*

The characteristics of Loomis's Mathematical Works, which we so highly prize, are fully sustained in his Natural Philosophy. I am glad to find at last a text-book on this subject so logically expressed, so complete, and so accurate."—N. B. WEBSTER, A. M., *President of Virginia Collegiate Institute, Portsmouth, Va.*

"This work of Professor Loomis claims to be a philosophical treatise. For clearness of style and fulness of illustration we have not met its equal. Adapted to the higher classes in academies and high schools, where mathematics have been pursued only through plane trigonometry, it is ready at all times to give a reason for a phenomenon, while the numerous references to the every-day occurrences of life make it exceedingly interesting to those who do not care for demonstrations. We hope it will crowd out of our schools the heterogeneous productions that school-boys have to stuff themselves with, and thus give thorough instruction, that will tend to develop some faculties besides memory."—S. J. C. SWEZEY, *Marion, Alabama.*

"Professor Loomis's Natural Philosophy combines conciseness with very great clearness of illustration, and is the best text-book of the kind that I have examined."—Prof. W. FERREL, *Nashville, Tenn.*

"Professor Loomis's treatise has my decided preference as a text-book on Natural Philosophy. It is simple and comprehensive in its style, and its principles are discussed with clearness and brevity."—J. W. FERREE, A. M., *Prof. of Mathematics, Dickinson Seminary, Pa.*

"Professor Loomis's Natural Philosophy is the best elementary work on this subject which I have hitherto seen. It is not only fitted for high schools, but is a perfect treasure for the private student.

Every subject on which it treats is pregnant with useful knowledge."—A. W. MASON, *Jeanesville Penn.*

"Professor Loomis's Natural Philosophy is the best text-book on that subject yet published. With all its value as a text-book, it combines a pleasing style, instructive examples and illustrations, and is quite a readable book for others besides school-boys."—B. W. McDONNOLD, *Professor in Bethel College, Tenn.*

"While this volume requires of the pupil nothing beyond a knowledge of the elementary mathematics, it develops the philosophy of all the leading principles of mechanics and physics. Its definitions are clear and concise, its statement of principles methodical and accurate. It embraces in a duodecimo volume of 350 pages a complete digest of what is usually spread over three or four times this space. Nor by this condensation is the study rendered repulsive, but exactly the reverse. We commend this work, without qualification, as one admirably adapted to interest the pupil, while it cultivates the habit of philosophical reflection and observation."—*New York Observer.*

"This volume is not only an admirable treatise for the use of the higher classes in schools, but is a most convenient compendium of the latest discoveries in Natural Philosophy for the instruction of adult readers."—*New York Commercial Advertiser.*

"This volume is an admirable compendium of the principles of Natural Philosophy; yet, strange to say, it is not dull, heavy, or incomprehensible, but states facts in terms that all can comprehend, and that few will be unable to remember."—*Baltimore Daily Times.*

"Among the numerous books of its class which have originated in this country, this volume has no superior in clearness, accuracy, and comprehensiveness."—*Boston Courier.*

"Professor Loomis's works on Arithmetic, Geometry, &c., are models of clear and precise style, comprehensive definitions, and accurate language. This work on Natural Philosophy is marked by the same excellences."—*Toronto Globe.*

"Ten pages of this book, which may easily be read by every mechanic in his winter evenings, will so fill his head, as to leave no room for idle thoughts."—*Newark Daily Advertiser.*

"The study of Professor Loomis's work will make solid thinkers, not smatterers. While popularizing science, he has not frittered away the solidity and accuracy which alone give it a title to that name."—*New York Daily News.*

### ALSO, PUBLISHED BY THE SAME AUTHOR,

#### A TREATISE ON ARITHMETIC,

Theoretical and Practical. Fifth Edition. 12mo. 352 pages. Sheep extra. 75 cents.

#### ELEMENTS OF ALGEBRA.

Designed for the Use of Beginners. Fourteenth Edition. 12mo. 281 pages. Sheep extra. 62½ cents.

#### A TREATISE ON ALGEBRA.

Twentieth Edition. 8vo. 359 pages. Sheep extra. \$ 1.00.

#### ELEMENTS OF GEOMETRY AND CONIC SECTIONS.

Eighteenth Edition. 8vo. 234 pages. Sheep extra. 75 cents.

#### TRIGONOMETRY AND TABLES.

Sixteenth Edition. 8vo. 360 pages. Sheep extra. \$ 1.50. The Trigonometry and Tables bound separately: The TRIGONOMETRY, \$ 1.00; TABLES, \$ 1.00.

#### ELEMENTS OF ANALYTICAL GEOMETRY,

And of the Differential and Integral Calculus. Thirteenth Edition. 8vo. 286 pages. Sheep extra. \$ 1.50.

#### INTRODUCTION TO PRACTICAL ASTRONOMY.

With a Collection of Astronomical Tables. Second Edition. 8vo. 497 pages. Sheep extra. \$ 1.50.

#### RECENT PROGRESS OF ASTRONOMY,

Especially in the United States. Revised Edition. 12mo. 396 pages. Muslin. \$ 1.00.

Also in Press, by the same Author,

#### ELEMENTS OF ASTRONOMY AND METEOROLOGY,

On the same plan as the "Elements of Natural Philosophy."

Published by HARPER & BROTHERS,  
Franklin Square, N. Y.

"SILLIMAN'S JOURNAL."

## THE AMERICAN JOURNAL OF SCIENCE AND ARTS,

(Two Volumes Annually, each of at least 450 pages, 8vo.)

PUBLISHED IN NUMBERS (ILLUSTRATED) OF 152 PAGES ON THE FIRST OF EVERY OTHER MONTH, VIZ.:—

JANUARY, MARCH, MAY, JULY, SEPTEMBER, & NOVEMBER,

AT NEW HAVEN, CONNECTICUT.

BY

B. SILLIMAN, JR., AND J. D. DANA.

\$5 PER ANNUM, IN ADVANCE.

The Journal is sent POST-PAID after the annual payment is received.

EDITED BY

PROFESSORS B. SILLIMAN, B. SILLIMAN, JR., AND PROF. JAMES DWIGHT DANA,  
NEW HAVEN,

IN CONNECTION WITH

Professor ASA GRAY, of Cambridge,

Professor LOUIS AGASSIZ, of Cambridge,

Dr. WOLCOTT GIBBS, of New York.

This work has now been established more than forty years, and is the ONLY JOURNAL of the kind in the United States. It is devoted to the general interests of PHYSICAL and CHEMICAL SCIENCE, GEOLOGY, MINERALOGY, NATURAL HISTORY, METEOROLOGY, GEOGRAPHY, and kindred departments of knowledge, and contains original papers, as well as abstracts of foreign discoveries, on all these topics.

Seventy-eight volumes have already been published,—FIFTY in the *first* and TWENTY-EIGHT in the *second* Series.

Subscribers receiving their copies direct from the Publishers remit their subscriptions to the Office of SILLIMAN'S JOURNAL, NEW HAVEN, CONN.

Most of the back volumes can be obtained of the Publishers.

All communications, remittances, &c., to be addressed to

**SILLIMAN & DANA,**

Office of Silliman's Journal of Science,

NEW HAVEN, CONN.

As an inducement to teachers and others, we offer to send this Journal for one year to any person who will send us the names of two new subscribers, accompanied by \$10, and for every additional new subscriber we will allow \$1 to be deducted from the remittance to accompany the name. Postage prepaid, in all cases, after the subscription for the current year is received.

### Notices.

A work which for nearly half a century has kept unflinchingly on the even tenor of its way, with a constantly growing reputation, has no need of culling "notices" from the Press. A single extract from a notice of the November number of 1859, from the Boston Courier and National Intelligencer, will show the estimation in which it is held.

"We are not aware that this number, whose contents we have briefly sketched, is richer than usual. We believe it steadily supports the character it has earned, of being, what good authorities have pronounced, the best scientific periodical in the English language which covers an equal, or nearly equal, field. Its editorial staff warrants its abilities. Besides its venerable founder, there is the younger Silliman, and Professor J. D. Dana, our most encyclopædic and ablest general naturalist, assisted by Agassiz, Gray, and Wolcott Gibbs in their several departments."

Confident as to the result of an impartial and intelligent examination, the Publishers will mail (post-paid) a copy of either of the works named below, to any teacher or school-officer remitting half the price.

A  
**NATURAL PHILOSOPHY,**  
FOR SCHOOLS AND COLLEGES.

By **G. P. QUACKENBOS, A.M.**

12mo. 450 pages. Price, \$1.00.

This book, which is illustrated with 335 fine engravings, is equally adapted to use with or without apparatus. Teachers will find it distinguished, in a remarkable degree, —

1. For clearness, fulness of explanation, and an original method of dealing with difficulties.
2. For its correction of numerous errors heretofore current in school Philosophies.
3. For its explanation of scientific principles as they appear in every-day life, and the practical application of these principles in problems presented for the pupils' solution.
4. The signal perspicuity of arrangement, everything being presented in its proper place.
5. For the interest with which it invests the subject. From the outset the student is fascinated.
6. For the embodiment of all recent discoveries. Instead of relying on the obsolete authorities that have furnished the matter for many of our popular school-books on this subject, the author has acquainted himself with the present state of science, and thus produced such a book as is demanded by the progressive spirit of the age.

Written by one of our most popular school-book authors, who admirably understands the art of adapting his instructions to the school-room, this work is acknowledged by the best educators in the land to be THE BOOK on Physical Science. Read the following extracts from letters, which might be extended through pages, did space allow.

"I have no hesitation in saying that, in my opinion, it is the best book on this subject with which I am acquainted. Such a text-book makes learning easy and teaching a pleasure." — *Geo. R. Perkins, LL.D., Author of Perkins's Mathematical Series.*

"It comes more fully up to the standard of a first-class text-book on Natural Philosophy for High Schools and Academies than any other work with which I am acquainted." — *Geo. A. Chase, Pres. of Brookville College, Brookville, Ind.*

"It has the same clearness, elegance, and finish that characterize your other productions." — *Tyler Lewis, LL.D., Prof. of Greek, Union College, Schenectady, N. Y.*

"It will soon become the favorite text-book of its kind in our Colleges and Schools; and this is a destiny which I honestly believe it richly deserves." — *Rev. W. C. Duncan, D.D., Late Prof. Latin and Greek, University of Louisiana, N. O.*

"I prefer it to any I have seen." — *W. E. Clifford, Prof. of Nat. Science, Mineral Point, Wis.*

"The book has no equal for the great mass of pupils in our Common Schools and Academies." — *A. J. Rickoff, Late City Supt. of Schools, Cincinnati, O.*

"It has been adopted by the Board of School Commissioners of this city, for use in all the Public Schools of the county. We regard it as far the best text-book on the subject now in use." — *P. A. Towne, General Princ. of the City Public Schools, Mobile, Ala.*

"I am led to regard the work as a model of what a school-book should be. None but a practical teacher could have so well understood, or so happily met, the wants of students." — *Isiah Peckham, Princ. of the High School, Newark, N. J.*

"It is the best book on the subject that I have seen. Its style is easy, its arrangement admirable, and its accuracy remarkable." — *J. G. Ralston, Princ. Oakland Fem. Inst. Norristown, Pa.*

"I regard it as better adapted to the general purposes of a School Philosophy

than any other text I have seen." — *Rev. R. H. Rivers, D.D., Pres. of Wesleyan Univ. Florence, Ala.*

"I fully concur in the opinion of Dr. R. H. Rivers, in reference to Quackenbos's Natural Philosophy." — *Rev. R. P. Ransom, A.M., Pres. Shelbyville (Tenn.) Univ.*

"In twenty-two years' experience in teaching, I have found no work more to my notion than this. It is so far ahead of the old text-books, that I am almost ashamed to own that I ever was an instructor in them." — *Rev. R. L. Abernathy, Princ. Rutherford Academies, Burke Co., N. C.*

"It is, in my opinion, unsurpassed as a text-book for all learners in this interesting department of study." — *Rev. Hubbard Winslow, D.D., Author of Intellectual Philosophy.*

"It is superior in many important points to the other elementary works on the same subject to which I have had access." — *Edward B. Smith, Princ. Piedmont Acad., Culpeper C. H., Va.*

"The Book is a perfect gem." — *John S. Hart, LL.D., Princ. High School, Philadelphia.*

"Its admirable illustrations, joined to its familiar application of principles and its practical problems, place it ahead of all competitors." — *C. W. Callender, Pres. Tennessee Female Coll., Franklin, Tenn.*

"It is decidedly the best Philosophy I have ever examined." — *Jesse W. Cornelius, Princ. of Academy, Cedarville, Ill.*

"It is a good book, and will meet with a large circulation." — *Rev. W. S. Barton, Editor Alabama Teacher, Montgomery, Ala.*

"We are using your Natural Philosophy in our school, and find it superior to any work we have ever used." — *Prof. J. W. Stewart, State Female College, Memphis, Tenn.*

"The clearness, simplicity, and accuracy, with which scientific principles are stated in this work, and the force with which they are illustrated, give it claim to a first place among the text-books on Philosophy." — *O. S. Dean, Princ. Susquehanna Collegiate Inst., Towanda, Pa.*

By the same Author.

ILLUSTRATED SCHOOL HISTORY OF THE UNITED STATES. 12mo. 473 pages. Price, \$1.00.

FIRST LESSONS IN COMPOSITION. 12mo. 182 pages. 50 cents.

ADVANCED COURSE OF COMPOSITION AND RHETORIC. 12mo. 450 pages. \$1.00.

Published by **D. APPLETON & CO.,**

346 and 348 Broadway, New York.







# ROBINSON'S COURSE OF MATHEMATICS.

BY HORATIO N. ROBINSON, LL.D.

<b>I. Robinson's Progressive Primary Arithmetic,</b>	<b>\$ 15</b>	<b>IX. Robinson's Key to University Algebra,</b>	<b>1 00</b>
<b>II. Robinson's Progressive Intellectual Arithmetic,</b>	<b>25</b>	<b>X. Robinson's Geometry and Trigonometry,</b>	<b>1 50</b>
<b>III. Robinson's Progressive Practical Arithmetic,</b>	<b>50</b>	<b>XI. Robinson's Surveying and Navigation,</b>	<b>1 50</b>
<b>IV. Robinson's Key to Practical Arithmetic,</b>	<b>50</b>	<b>XII. Robinson's Analytical Geometry and Conic Sections,</b>	<b>1 50</b>
<b>V. Robinson's Progressive Higher Arithmetic,</b>	<b>75</b>	<b>XIII. Robinson's Differential and Integral Calculus,</b>	<b>1 50</b>
<b>VI. Robinson's New Elementary Algebra,</b>	<b>75</b>	<b>XIV. Robinson's Elementary Astronomy,</b>	<b>75</b>
<b>VII. Robinson's Key to Elementary Algebra,</b>	<b>60</b>	<b>XV. Robinson's University Astronomy,</b>	<b>1 75</b>
<b>VIII. Robinson's University Algebra,</b>	<b>1 25</b>	<b>XVI. Robinson's Mathematical Operations,</b>	<b>2 25</b>
		<b>XVII. Robinson's Key to Algebra, Geometry, &amp;c., (In Press.)</b>	

For extent of research, facility and aptness of illustration, and practical usefulness, the author of this series is surpassed by no mathematical writer in this country. This Series has been cordially recommended by many of the most eminent Mathematicians in the country. Among them,—

**Hon. W. C. Larrabee**, Superintendent Public Instruction, Indiana.  
**Hon. Jona. Tenny**, Secretary Board of Education, N. H.  
**Hon. Lyman C. Draper**, Superintendent Public Instruction, Wisconsin.  
**Hon. S. H. Carpenter**, Assistant Superintendent, Wisconsin.  
**Prof. Dewey**, Rochester University.  
**Prof. McFarland**, Miami University.  
**Prof. Thompson**, Hanover College.  
**Prof. Wilson**, Hobart Free College.

**Prof. Stone**, Iowa University.  
**Prof. Beattie**, Wesleyan University.  
**President Rawlins**, Asbury Female College.  
**Prof. Standish**, Knox College.  
**Prof. Hudnot**, Lind University, Ill.  
**Prof. H. Pomeroy**, Lawrence University, Wis.  
**Rev. Dr. Conrad**, Cooper Female Institute.  
**President D. Burbank, LL. D.**, Brockport Collegiate Institute.  
**Hon. Henry Kiddle**, Assistant Superintendent, New York city.

## NEW RECOMMENDATIONS.

In the January and March No.'s of **IVISON & PHINNEY's Educational News**, a list of recommendations of **ROBINSON'S WORKS** was given, including the testimonials of several Professors of Mathematics, and of more than **TWENTY COUNTY COMMISSIONERS** of the State of New York, and of several Teachers' Associations. Since that time, and within three months, the publishers have been favored with a mass of Testimonials from most important sources, which can only be referred to in this place, and which furnish the highest proofs of the great value and adaptation of the works. The following names are selected from hundreds of eminent Teachers who have recently given cordial and discriminating commendations.

**Prof. D. Wood**, Prof. Engineering and Physics, University of Michigan.  
**Prof. James C. Watson**, Assistant in Observatory, University of Michigan.  
**Prof. S. S. Greene**, Brown University.  
**Prof. J. W. Patterson**, Dartmouth College.  
**Prof. T. Mc. Ballentine**, Prof. of Mathematics, Cumberland College, Ky.  
**W. H. Seavey, A. M.**, Principal Girls' High and Normal School, Boston.  
**Prof. Dascom Greene**, Prof. of Mathematics, Troy Polytechnic Institute.  
**Prof. J. W. Sterling, A. M.**, Prof. of Mathematics, Wisconsin University.  
**Prof. C. Tracy**, Prof. of Mathematics, State Agricultural College, Michigan.  
**Prof. C. T. Farrar**, Elmira Female College.  
**Prof. S. I. Fowler**, Prof. of Mathematics, Hillsdale College, Michigan.  
**D. B. Paine**, Byhalia Academy, Mississippi.  
**Prof. W. A. Clark**, Principal Genesee Wesleyan Seminary.  
**Prof. W. S. Rogers**, Alfred Academy, New York.  
**Prof. N. W. Butts**, Principal Medina Academy, New York.  
**O. S. Throop**, President Genesee County Teachers' Association.

**Prof. C. Z. Case**, Principal Red Creek Academy, New York.  
**Z. L. Parker**, Principal Union School, Corning, New York.  
**Prof. W. L. Harvey**, Lagrange, Mo.  
**Prof. A. C. Richmond**, Cotton Hill, Georgia.  
**Prof. J. S. French**, London Bridge, New Hampshire.  
**James Atwater, Esq.**, Superintendent Schools, Lockport, N. Y.  
**Prof. E. A. Charlton, A. M.**, Principal Union School, Lockport, N. Y.  
**Prof. C. W. Sanders, A. M.**, New York.  
**Prof. E. Chamberlain**, Elkhorn Academy, Wisconsin.  
**J. K. Comstock**, West Onondaga, N. Y.  
**Prof. W. L. Hall, A. M.**, Principal Union School, Bath, N. Y.  
**E. B. Hall**, Union School, Fairport, N. Y.  
**Prof. Rufus Sheldon**, Principal Auburn Academy, N. Y.  
**Prof. Charles Fairman**, Prin. Yates Academy, N. Y.  
**B. C. Rude**, Principal Union School, Newark, N. Y.  
**Prof. D. J. Pratt, A. M.**, Principal Academy, Fredonia, N. Y.  
**T. K. McArthur**, North Sharon, Pa.  
**Prof. D. Brown, A. M.**, Cockrum, Miss.  
**Prof. J. B. Snow**, Principal Union School, Lyons, N. Y.  
**Prof. T. Baldwin**, Indiana Normal Institute, Burnettsville, Ind.

For 25 SOLID PAGES OF TESTIMONIALS from practical and successful educators, see **IVISON, PHINNEY, & Co.'s DESCRIPTIVE CATALOGUE**, and "NEWS."

Single copies of such of these books as are used in classes sent at half price to Teachers for examination with a view to introduction. Liberal terms for first supplies.

**IVISON, PHINNEY, & Co.,**

Publishers, 48 and 50 Walker Street, New York.

# DISTINGUISHED EDUCATORS

## WHO HAVE COMMENDED WORKS IN THE AMERICAN EDUCATIONAL SERIES

— OF —

### SCHOOL AND COLLEGE TEXT BOOKS, ETC.,

PUBLISHED BY

## IVISON, PHINNEY, & CO.,

40 & 50 Walker St., New York.

- |  |  |  |
|--|--|--|
| <p>Pres. NOTT, D.D., LL.D., Union College.<br/>           Pres. HOPKINS, D.D., Williams College.<br/>           Pres. DAY, D.D., LL.D., Yale Coll.<br/>           Pres. HITCHCOCK, D.D., LL.D., Amherst College.<br/>           Pres. SEARS, D.D., Brown University.<br/>           Pres. FRELINGHUYSEN, LL.D., Rutgers College.<br/>           Pres. ANDREWS, D.D., Marietta College.<br/>           Pres. ANDERSON, D.D., Rochester University.<br/>           Pres. HORACE MANN, LL.D., Antioch College.<br/>           Pres. ALDEN, D.D., Jefferson Coll.<br/>           Pres. COLLINS, D.D., Dickinson College.<br/>           Pres. ANDREWS, D.D., Gambier College.<br/>           Prof. A. D. BACHE, Washington, D. C.<br/>           Prof. G. P. BOND, Harvard College.<br/>           Dr. B. A. GOULD, Jr., Albany.<br/>           Prof. CHARLES W. HACKLEY, Columbia College, N. Y.<br/>           Rev. THOMAS HILL, Pres. Antioch College.<br/>           Prof. E. OTIS KENDALL, University of Pennsylvania.<br/>           Prof. BENJAMIN PEIRCE, Harvard University.<br/>           Prof. PARSONS, LL.D., Harvard University.<br/>           Prof. SHEDD, D.D., Andover Seminary.<br/>           Prof. HAYES, M.D., State Assayer, Massachusetts.</p> | <p>Prof. SANBORN, Dartmouth Coll.<br/>           Prof. PORTER, D.D., Yale College.<br/>           Prof. DANA, LL.D., Yale College.<br/>           Prof. TUCKERMAN, Amherst Coll.<br/>           Prof. CLARK, Amherst College.<br/>           Prof. PERRY, Williams College.<br/>           Prof. CHADBOURNE, Williams College.<br/>           Prof. NORTH, Hamilton College.<br/>           Prof. HOPKINS, Genesee College.<br/>           Prof. SCHMIDT, Columbia College.<br/>           Pres. WEBSTER, LL.D., New York Free Academy.<br/>           Pres. THOMPSON, D.D., Hanover College.<br/>           Pres. GERHART, D.D., Franklin and Marshall College.<br/>           Pres. BAUGHER, D.D., Pennsylvania College.<br/>           Pres. HENRY, Smithsonian Institute.<br/>           Prof. AGASSIZ, Harvard University.<br/>           Prof. PEARSON, Union College.<br/>           Prof. LEWIS, LL.D., Union College.<br/>           Prof. SILJMAN, LL.D., Yale Coll.<br/>           Prof. GUYOT, LL.D., Princeton College.<br/>           Prof. SCHANCK, M.D., Princeton College.<br/>           Prof. TORREY, LL.D., United States Assay Office.<br/>           Prof. DEWEY, LL.D., Rochester University.<br/>           Prof. McFARLANE, Miami Univ.<br/>           Prof. GREENE, C. E., Troy Polytechnic Institute.<br/>           Prof. STONE, Iowa University.<br/>           Prof. POMEROY, C. E., Lawrence University.<br/>           Prof. J. W. STERLING, Wisconsin University.</p> | <p>Prof. J. O. HUDNUT, C. E., Lind University.<br/>           Prof. STANDISH, Knox College.<br/>           Prof. BEATTIE, Wesleyan Univ.<br/>           Prof. HOVEY, Normal University, Illinois.<br/>           Prof. WINCHELL, Michigan Univ.<br/>           Prof. MADLIN, University North Carolina.<br/>           Prof. SHIFF, University North Carolina.<br/>           Prof. J. S. HART, LL.D., Phila.<br/>           Prof. BARNES, U. S. Naval Acad.<br/>           Prof. EVANS, Marietta College.<br/>           Prof. HILLMAN, Dickinson College.<br/>           Prof. SHERWIN, City Normal School, Boston.<br/>           Prof. SEAVEY, Girls' Norm. School, Boston.<br/>           Prof. PHELPS, New Jersey Normal School.<br/>           Prof. BEAN, Normal School, Bridgeton, Maine.<br/>           Prof. WEBSTER, Portsmouth, Va.<br/>           Prof. HADLEY, Wabash College, Iowa.<br/>           Prof. LINDSLEY, Nashville, Tenn.<br/>           Prof. PRICE, University Missouri.<br/>           Prof. SWALLOW, Univ. Missouri.<br/>           Prof. HENPHILL, University South Carolina.<br/>           Prof. ROCKWELL, Davidson Coll.<br/>           Prof. CRAVEN, North Carolina Normal College.<br/>           Pres. CUMMINGS, D.D., Asheville, North Carolina.<br/>           Prof. HARRIS, Lagrange College, Louisiana.<br/>           And many STATE SUPERINTENDENTS and others.</p> |
|--|--|--|

### IVISON, PHINNEY, & CO.'S NEW DESCRIPTIVE CATALOGUE.

**168 Pages, 8vo.**

Owing to the great number of important and influential Testimonials which have been volunteered since the issue of their former CATALOGUE, the Publishers have issued a

#### NEW DESCRIPTIVE CATALOGUE OF 168 PAGES,

Filled with extracts from thousands of Testimonials from Teachers and the Press in all parts of the Union, which will be sent, pre-paid, on application. SINGLE COPIES of any of the books of the AMERICAN EDUCATIONAL SERIES, that are USED IN CLASSES, will be sent at HALF-PRICE to TEACHERS for Examination, with a view to Introduction. Any books sent, pre-paid, to any address, on receipt of retail price.

*SPECIALLY FAVORABLE TERMS FOR FIRST INTRODUCTION.*

Published by **IVISON, PHINNEY, & CO.,**

48 and 50 Walker Street, NEW YORK.

SOLD BY BOOKSELLERS GENERALLY.